


Biologically guided flap stability: the role of flap thickness including periosteum retention on the performance of the coronally advanced flap—A double-blind randomized clinical trial

Marco Clementini¹  | Nicola Discepoli² | Carlotta Danesi¹ | Massimo de Sanctis¹

¹Department of Periodontology, Università Vita-Salute San Raffaele, Milan, Italy

²Department of Medical Biotechnologies, Unit of Periodontics, Università degli Studi di Siena, Siena, Italy

Correspondence

Marco Clementini, Department of Periodontology, Università Vita-Salute San Raffaele, Milan, Italy.
Email: mclementini@me.com

Abstract

Aim: To evaluate the possible benefit on wound healing and flap stability of periosteum inclusion, comparing a “split–full–split” thickness flap elevation versus a “split” thickness approach performed during CAF for the treatment of isolated-type gingival recessions in the upper jaw.

Material and Methods: Forty patients were randomized, 20 were treated with “split–full–split” (test group) and 20 with a “split” approach (control group). Analysed parameters at 1 year were CRC, percentage of recession coverage (RC), keratinized tissue (KT) gain and patient-related outcome measurements.

Results: After 12 months, CRC was 80% in the test group and 35% in the control group. Percentages of RC and KT gain were higher in the test group, and a significant association between CRC and the thickness of the flap after elevation was found. Patient-related outcomes measurements were better for the test group.

Conclusions: Flap thickness preservation and the presence of the periosteum in part of the flap may play a fundamental role in obtaining CRC.

KEYWORDS

complete root coverage, coronally advanced flap, flap elevation, flap thickness, gingival recession

1 | INTRODUCTION

Treatment of buccal gingival recession (GR) is the common clinical requirement from patients who are mainly concerned about aesthetics. Noteworthy are also requests linked to root sensitivity, difficulty in oral hygiene procedures, presence of root caries and noncarious cervical lesions (Tonetti & Jepsen, 2014). GR defects, when left untreated, do not improve spontaneously and may progress towards increased recession depth (RD) and clinical attachment loss which increase the patient's aesthetic concern and the clinical discomfort

due to augmented dental hypersensitivity (Chambrone & Tatakis, 2016).

Complete root coverage (CRC) can be considered the primary clinical outcome (Chambrone & Tatakis, 2015) and selecting the surgical technique depends mainly on the local anatomical characteristics and on the patient's demands (de Sanctis & Clementini, 2014).

In patients with a residual amount of keratinized tissue apical to the recession defect, the coronal advanced flap (CAF) may be recommended. This surgical technique results in optimal root coverage, good colour blending of the treated area with respect to adjacent

soft tissues and a complete recovery of the original (presurgical) soft tissue marginal morphology (de Sanctis & Zucchelli, 2007). Furthermore, postoperative morbidity is reduced to a single area of surgical intervention and the overall chair time is limited.

When utilizing CAF technique, critical factors in CRC have been described in the literature. Flap positioning coronal to the CEJ (Pini Prato et al., 2005) and a tension-free flap design (Pini Prato et al., 2000) are among the most important ones. Moreover, flap thickness has been shown to influence the clinical outcomes of CAF procedure (Baldi et al., 1999).

Coronally advanced flap has been widely validated by the literature for the treatment of single recession defects (Cairo, Nieri, & Pagliaro, 2014) and, currently, different flap designs and technical modifications are available to clinicians (Allen & Miller, 1989; Bernimoulin, Luscher, & Muhlemann, 1975; Norberg, 1926; Pini Prato et al., 1992).

de Sanctis and Zucchelli (2007) have introduced the "split-full-split" flap elevation modality. According to the authors, the modulation of flap thickness, produced by the inclusion of periosteum in the central area, increases flap thickness in the portion of the flap residing over the previously exposed avascular root surface. This, in turn, would give better stability to the flap. However, the partial-thickness flap approach used to be commonly performed and taught in the clinical practice, and it is validated in the literature (Allen & Miller, 1989; da Silva, Joly, de Lima, & Tatakis, 2004).

To date, evidence is still lacking on the influence of including the periosteum in the flap when compared with a split thickness approach in obtaining a CRC.

Thus, the aim of this double-blind, controlled and randomized clinical trial was to evaluate the possible benefit on wound healing and flap stability of periosteum inclusion comparing a "split-full-split" flap elevation versus a "split" thickness approach when CAF is performed for the treatment of isolated-type gingival recessions in the upper jaw.

2 | MATERIALS AND METHODS

2.1 | Study design

This investigation was a parallel, randomized, single-centre clinical trial with blinded outcome assessment on the treatment of single maxillary gingival recession defects according to the CONSORT statement (<http://www.consort-statement.org/>). The study was registered on ClinicalTrial.org (ID: NCT03417232). Two different treatment modalities were compared: the CAF with a "split-full-split" thickness (SFPT) flap elevation (test group) and the CAF with a partial-thickness (PT) flap elevation (control group). The flow chart of the study is presented in Figure 1.

2.2 | Population

Before any therapy was accomplished, in full accordance with the ethical principles of the Declaration of Helsinki on experimentation involving human subjects, as revised in 2000, the protocol was approved by the University Ethical Board (Ref. CAF0001 23.04.13)

CLINICAL RELEVANCE

Scientific rationale for the study: There is a lack of evidence on the influence of including or not the periosteum in the flap during elevation of a coronally advanced flap for obtaining a CRC.

Principal findings: Frequency of CRC and percentages of RC showed a superior performance for the "split-full-split" thickness (test) than for the "split" thickness (control) elevation. Moreover, patient-related outcome measurements achieved in the test group were better than the control group.

Practical implications: Flap thickness preservation and inclusion of the periosteum in the flap during elevation of a coronally advanced flap may play a role in the achievement of better recession reduction outcomes and patient-related outcomes.

at University of Siena, Italy. Patients agreed to participate in the study by signing a written informed consent according to the above-mentioned principles. The participants were selected on a consecutive basis among patients of the Department of Periodontology at the University of Siena, Italy, between April 2013 and April 2015.

2.3 | Inclusion criteria

In order to be included in the current clinical investigation, patients had to meet the following criteria:

- age >18 years,
- no systemic diseases or pregnancy,
- smoking ≤10 cigarettes/day,
- full-mouth plaque score and full-mouth bleeding score ≤20%,
- presence of at least one Miller class I isolated recession defect (Miller, 1985) in the upper jaw and at least 2 mm of keratinized tissue apical to the recession,
- recession depth (RD) equal to or greater than 2 mm,
- identifiable cemento-enamel junction (CEJ),
- vital teeth, free from caries or prosthetic crown,
- no history of periodontal surgery at experimental sites.

2.4 | Randomization, allocation concealment and blinding

A blocked randomization (for gender and smoking status) was performed using a computer software by someone not involved in other aspects of the study. Allocation concealment was performed by opaque sealed envelopes that were opened after designing the flap over the soft tissues. The surgeon was not blind to treatments. Examiners, patients and statisticians were blind to procedures.




FIGURE 1 CONSORT flow chart of the study

2.5 | Clinical assessment

The following parameters were recorded with a periodontal probe (PCP UNC 15, Hu-Friedy) at baseline and 12 months:

- full-mouth plaque score (FMPS) (O'Leary, Drake, & Naylor, 1972) and presence/absence of visible plaque.
- full-mouth bleeding score (FMBS) (Muhlemann & Son, 1971) and presence/absence of bleeding on probing at study tooth site.
- recession depth (RD), measured from the cemento-enamel junction (CEJ) to the most apical extension of the gingival margin.
- probing depth (PD), measured from the gingival margin to the bottom of the gingival sulcus.
- clinical attachment level (CAL), measured from the CEJ to the bottom of the gingival sulcus.
- keratinized tissue height (KTH), measured from the most coronal extension of gingival margin to the mucogingival line.

2.6 | Other parameters were recorded, as

- Dentin hypersensitivity (DH), assessed by 10 s air spray applied to the exposed buccal cervical area,

- Gingival thickness (GT) at baseline, determined by inserting a needle with a silicon stop, perpendicular to the tissue surface 3 mm apical to the gingival margin. After reaching the hard surface, the silicon stop was slid and placed in contact with the soft tissue (Zucchelli et al., 2010). After removing the needle, the distance between the tip of our needle and the silicon stop was measured with a calliper accurate to the nearest 0.1 mm.
- Flap thickness (FT), recorded intrasurgically after elevation 3 mm apical to the gingival margin, using a Iwanson gauge, modified (without the spring) in order to avoid excessive pressure on the soft tissue (Baldi et al., 1999).

2.7 | Patient-centred outcomes

A visual analogue scale (VAS) was used to assess:

- Postoperative discomfort, bleeding and swelling, at 7 days after surgery (0 = no discomfort at all and 10 = extreme discomfort).
- Aesthetic evaluation, in terms of root coverage and colour match, at 12 months (0 = bad aesthetic and 10 = good aesthetic).
- Satisfaction, in terms of overall satisfaction to the treatment, at 12 months (0 = no satisfaction at all and 10 = high satisfaction).

2.8 | Investigator training

All surgeries were performed by the same operator (MC), and all clinical measurements were performed by two blinded examiners (N.D, C.D.). They were calibrated by measuring twice, 24 hr apart, the PD and the RD of three patients not included in the study. Kappa statistic was used to determine inter- and intraexaminer reproducibility.

2.9 | Pretreatment: modification of oral hygiene habits

After the screening examination, all subjects received a session of prophylaxis including proper oral hygiene instructions (OHI), scaling and professional tooth cleaning, by means of a rubber cup and a low abrasive polishing paste. In the presence of recession-type defects, a "coronally directed" roll technique was suggested to minimize trauma at the gingival margin. Surgical treatment of recession defects was not performed until all patients displayed an adequate standard of supragingival plaque control (FMPS < 20%), as well as low tissue inflammation (FMBS < 20%).

2.10 | Surgical technique

The design of the flap consisted of two horizontal beveled incisions (3 mm in length), mesial and distal to the recession defect, and two slightly oblique beveled incisions. In the test site (SFST), the resulting trapezoidal-shaped flap was elevated with a split–full–split approach in the coronal–apical direction (de Sanctis & Zucchelli, 2007). The surgical papillae comprised between the horizontal incisions and the sulcular area apical to the recession was both elevated split thickness. Conversely, the central portion of the flap apical to the recession was elevated full thickness by the use of a small periosteum elevator inserted into the probable sulcus up to exposing 3–4 mm of bone apical to the bone dehiscence (Figure 2). In the control

group (PT), the flap was fully elevated with a split thickness approach, inserting the blade (15c) of the knife into the sulcus and continuing dissection with the blade to the mucogingival line (Allen & Miller, 1989; da Silva et al., 2004) (Figure 3). In both groups, the incision was terminated apically to the mucogingival line by means of a split thickness elevation to free the flap from muscle tension, keeping the blade (15c) parallel to the external mucosal surface. When the margin of the flap reached passively a level coronal to the CEJ of the tooth affected by recession, the coronal mobilization was considered adequate. Only the portion of root surface relative to loss of clinical attachment was mechanically treated utilizing curettes, and the application of EDTA for two minutes followed by a rinse of physiologic solution. The flap was positioned 1 mm coronal to the CEJ. The suture of the flap started with two interrupted periosteal sutures performed at the most apical extension of the vertical releasing incisions; afterwards, it proceeded coronally with other interrupted sutures, each of them directed, from the flap to the adjacent buccal soft tissue, in the apical–coronal direction. The last sling suture allowed for the stabilization of the surgical papillae over the interdental connective tissue bed and allowed for a precise adaptation of the flap margin over the underlying convexity of the crown.

2.11 | Postsurgical infection control

Patients were instructed to rinse with chlorhexidine solution (0.12%) twice a day for 1 min. Fourteen days after surgery, sutures were removed and the patient was prompted to rinse with chlorhexidine for 2 more weeks to maintain plaque control in the treated area. After this period, patients were again instructed to a mechanical cleaning of the treated region using a postsurgical toothbrush. All patients were recalled for control appointments at 2, 4 and 6 weeks after suture removal and, subsequently, once every 3 months until 1-year follow-up.




FIGURE 2 Split–full–split approach (test group) for the treatment of a localized gingival recession




FIGURE 3 Split approach (control group) for the treatment of a localized gingival recession

2.12 | Sample size determination

The sample dimension was calculated using $\alpha = 0.05$ and the power ($1-\beta$) of 90%. The minimum expected effect size was established on a difference of 515%, from 37% (the median of CRC for CAF alone of included studies in the systematic review by Cairo et al., 2014) to 88% (the higher CRC for CAF alone of included studies in the systematic review by Cairo et al., 2014). Calculations were performed using Stata 15 IC (Stata corp.). On the basis of these data, the needed number of patients to be enrolled in this study was 20 for the test group and 20 for the control. However, the number of patients was increased of 20% each arm considering the possibility of drop out.

2.13 | Data analysis

Descriptive statistics were expressed as mean, standard deviation (SD) and confidence of interval at 95% (CI: 95%). Percentages of RC and the achievement of CRC at 12 months were calculated.

The primary outcome was determined by the number of patients with recession defects that obtained CRC. Secondary outcomes were the percentages of RC, change in RD, KTH and DH, patient's discomfort (VAS discomfort), patient's preference in terms of aesthetic result (VAS aesthetic) and patient's satisfaction (VAS satisfaction).

Student's *t* test was used to evaluate the differences between groups regarding RD, PD, GT, FT, percentage of RC, KTH, DH, VAS discomfort, VAS aesthetic and VAS satisfaction. Paired *t* test was used to evaluate the differences between baseline and 1-year follow-up of RD, PD, KTH, DH. The Pearson's chi-square test was used to compare the two groups regarding CRC. A logistic regression model was fitted to relate CRC as dependent variable and intraoperative FT as independent variable ($p < 0.05$).

3 | RESULTS

3.1 | Study population

A total of 50 patients were included in the study: Three patients declined to participate and three patients did not meet the inclusion criteria. Forty-four patients completed the study procedures. Since

four patients dropped out during follow-up, 40 patients were finally analysed (CONSORT flow chart, Figure 1).

During the study period, all patients presented a good standard of supragingival plaque control. Experimental (test and control) sites did not show BOP or visible plaque.

No differences were present between groups regarding patient age (test: 38.4 ± 9 years [CI 95%: 30.4–45.3]; control: 36.4 ± 12 years [CI 95%: 29.6–43.2]; $p > 0.5$), gender (test: 15 females; control: 14 females; $p > 0.5$) tooth type distribution (test: 2 incisors, 12 canines, 8 premolars; control: 3 incisors, 10 canines, 9 premolars; $p > 0.5$) and smoking status (test: 10 smokers; control: 12 smokers; $p > 0.5$). (Table 1).

3.2 | Clinical parameters

Kappa score used to determine intra- and interexaminer reproducibility was 0.88 and 0.82, respectively.

The descriptive statistics for the clinical parameters measured at baseline and 1 year after surgery for both groups are shown in Table 2.

At baseline, there was no significant difference between both groups regarding any of the clinical parameters. RD was 2.47 ± 0.9 mm (CI 95%: 2–3) for the test group and 2.33 ± 0.9 mm (CI 95%: 1.9–2.8) or the control group ($p > 0.5$), and PD was 1.4 ± 0.7 mm (CI 95%: 1–2) for the test group and 1.3 ± 0.8 mm (CI 95%: 1–2) for the control group ($p > 0.5$). KTH was 2.5 ± 0.7 mm (CI 95%: 2–1.3) for the test group and 2.6 ± 1 mm. (CI 95%: 2–3.2) for the control group ($p > 0.5$).

Gingival tissue thickness at baseline was 0.9 ± 0.2 mm (CI 95%: 0.8–1.1) for the test group and 0.9 ± 0.1 mm (CI 95%: 0.8–1.1) for the control group ($p > 0.5$). After elevation, flap thickness was significantly different ($p < 0.01$) between the test group (0.93 ± 0.1 mm. [CI 95%: 0.8–1.1]) and the control group (0.46 ± 0.1 mm. [CI 95%: 0.4–0.5]).

After 12 months, PD did not change in both groups ($p > 0.5$) whereas RD significantly decreased for both groups ($p < 0.01$) compared to baseline measurements. A significant between-groups difference was found at 1 year, with the test group

TABLE 1 Study Population

	"split-full-split" (N = 22)	"split" (N = 22)
Age (Confidence interval 95%)	38.4 ± 9 years (30.4–45.3)	36.4 ± 12.2 years (29.6–43.2)
Gender	15 females 7 males	14 females 8 males
Tooth type	2 incisors 12 canines 8 premolars	3 incisors 10 canines 9 premolars
Smokers	10	12

TABLE 2 Clinical parameters

	"split-full-split"(N = 20)	"split"(N = 20)
RD (mm)		
Baseline	2.47 ± 0.9(2–3)	2.33 ± 0.9(1.9–2.8)
12 months	0.23 ± 0.5* [#] (0–0.5)	0.76 ± 0.7* ^x (0.3–1.2)
PD (mm)		
Baseline	1.4 ± 0.7 (1–2)	1.3 ± 0.8(1–2)
12 months	1.3 ± 0.8(1–2)	1.2 ± 0.4(1–2)
KT (mm)		
Baseline	2.5 ± 0.7(2.1–3)	2.6 ± 1(2–3.2)
12 months	2.7 ± 0.8(1.8–2.8)	2 ± 0.8*(1.6–2.4)
GT (mm)		
Presurgical	0.9 ± 0.2(0.8–1.1)	0.9 ± 0.1(0.8–1.1)
FT (mm)		
Intrasurgical	0.93 ± 0.1** ^(0.8–1.1)	0.46 ± 0.1(0.4–0.5)
RC (%)		
12 months	92.3 ± 16.6** ^(83.1–100)	72.5 ± 22.4(60–84.9)
CRC (%)		
12 months	80% (16/20)**	35% (7/20)

Notes. (Confidence interval 95%).

*Statistically significant difference within two groups, **Statistically significant difference between two groups.

presenting higher RD reduction ($p < 0.5$), higher percentages of CRC (80% for test group and 35% for control group, with $p < 0.01$) and higher percentages of RC (92.3 ± 16.6 [CI 95%: 83.1–100] for test group and 72.5 ± 22.4 [C.I 95%: 60–84.9] for control group, with $p < 0.01$). KTH slight decreased in the control group ($p > 0.5$) while it slightly increased in the test group ($p = 0.5$).

Logistic regression analysis showed a significant association between CRC and the thickness of the flap resulted after elevation ($p < 0.01$).

3.3 | Patient-centred outcomes

The descriptive statistics for patient-centred outcome are shown in Table 3.

Hypersensitivity was present at baseline in 11/20 (55%) of sites in the test group and 12/20 (60%) of sites in the control group ($p > 0.5$) while at 1 year it was present in 0/20 (0%) of sites in the test group and 4/20 (20%) of sites in the control group ($p < 0.5$).

VAS discomfort during the first week was 2.3 ± 2.5 (CI 95%: 0.9–3.7) for the test group and 5.5 ± 2.4 (CI 95%: 3.8–7.3) for the control group, showing a statistically significant difference between groups ($p < 0.01$).

VAS aesthetic and satisfaction at 12 months were, respectively, 8.80 ± 1 (CI 95%: 8.2–9.3) and 8.47 ± 1.1 (CI 95%: 8.2–9.3) or the test group and 8.20 ± 1.6 (CI 95%: 7.8–9.3) and 8.20 ± 1.6 (CI 95%: 7.8–9) for the control group, with no statistically difference ($p = 0.2$ for VAS aesthetic and $p = 0.7$ for VAS satisfaction). Considering a range from 5 (neutral) to 10 (very good) of the VAS scale, a significant difference was found in favour of the test group as for VAS satisfaction ($p < 0.5$) but not for VAS aesthetic ($p = 0.05$).

4 | DISCUSSION

The present parallel, double-blinded, randomized clinical trial compared clinical and patient-centred outcomes between coronally advanced flaps (CAF) for the treatment of Miller's type I and II gingival recessions. Periosteum retention ("split-full-split" approach, test group) provides superior clinical results in terms of CRC (main outcome variable) and RC; hypersensitivity and postsurgical discomfort were also significantly improved.

The main outcome variable of the current investigation was the CRC. The frequency of CRC achieved by the test group was twice as high as that of the control group. Also for root coverage (RC), as a secondary outcome variable, the results obtained show a superior performance for the SFST (test) than for the ST (control).

TABLE 3 Patient-centred outcomes

	"split-full-split"(N = 20)	"split"(N = 20)
Hypersensitivity		
Baseline	60% (12/20)	55% (11/20)
12 months	0% (0/20)* ^{***}	20% (4/20)*
VAS discomfort		
7 days	2.3 ± 2.5 ** ^(0.9–3.7)	5.5 ± 2.4 (3.8–7.3)
VAS satisfaction		
12 months	8.47 ± 1.1 ** ^(7.8–9)	8.20 ± 1.6 (7.89.3)
VAS aesthetic		
12 months	8.80 ± 1 (8.2–9.3)	8.20 ± 1.6 (7.3–9)

Notes. (Confidence interval 95%).

*Statistically significant difference within two groups, **statistically significant difference between two groups, ***Statistically significant difference when VAS range was considered from 5 (neutral) to 10 (very good).

The marked clinical differences obtained between the two experimental groups seem to confirm the biological rationale that we are trying to test, so that the periosteum included within the flap and the use of the elevator in detaching the sulcular area may contribute to a better flap stability over the avascular root surface and a consequent better clinical result.

The surgical technique that detaches the marginal tissue by means of periosteum elevator instead of a surgical blade may prevent the reduction in the marginal thickness by avoiding involuntary microwounds of the marginal area and by maintaining the entire structure of the sulcular area. The major thickness of the marginal area can have a role in absorbing or deflecting wound-rupturing forces that otherwise would have been transmitted to the fibrin clot at the root surface—mucogingival flap interface.

From a biological standpoint, the role played by the periosteum seems to be crucial. Actually, during the very early phases of healing, the stability of a new fragile matrix (blood clot) is challenged by mechanical forces acting on the wound margins (Kon, Novaes, & Ruben, 1969; Laurens et al., 2006; Wikesjö & Nilvéus, 1990). The split-full-split flap modulation guarantees several biological conditions that may help a new connective tissue formation during these early phases of wound healing. *In vivo* preclinical studies have shown how the clot stability and its adhesion to the root surface positively influence the formation of a new attachment instead of a long junctional epithelium (Haney, Nilvéus, Mc Millan, & Wikesjö, 1993; Wikesjö, Claffey, & Egelberg, 1991; Wikesjö, Claffey, Nilvéus, & Egelberg, 1991). In this study, the intrinsic stiffness of the periosteum included within the flap has probably maintained the maximum thickness of the most critical area, that is the avascular root surface. Furthermore, the periosteum, with its high vascularization, provides an important source of cells (fibroblasts and macrophages above all) and growth factors (transforming growth factors beta family). Over the previously denuded root surface, the periosteum indeed can accelerate the phase of new tissue formation, acting as a source of endothelial cells (Potente, Gerhardt, & Carmeliet, 2011) and fibroblasts (Grieb, Steffens, Pallua, Bernhagen, & Bucala, 2011); the availability of these cell type in the periodontal wound is crucial to speed up the transition from a catabolic to an anabolic phase and, hence, the maturation/stabilization of the blood clot (Grieb et al., 2011).

Our data agree with those published in a recent systematic review in which a range of 34.2%–96.6% for RC and 7.7%–88% for CRC was reported when a single recession was treated by a CAF (Cairo et al., 2014). Results from meta-analysis showed that CAF plus connective tissue graft (CAF+CTG) or CAF plus enamel matrix derivative (CAF+EMD) was more effective than CAF alone in terms of CRC and RC. Another systematic review (Chambrone & Tatakis, 2015) confirmed the superiority of CAF plus SCTG to provide the best clinical outcomes (CRC, RC) in the treatment of single maxillary recession. Interestingly, the test group in the current investigation has achieved CRC with a frequency quite similar to those found in the systematic reviews by the groups CAF plus SCTG. It is well known that the presence of a connective tissue graft beneath a surgical flap leads to higher percentages of root coverage because it

reduces the space for a better blood clot stability and maturation. These findings suggest that a modulate flap thickness (“split–full–split”) may partially mimic the beneficial effects on flap stability that are obtained when a connective tissue graft is added.

According to the modification of the technique, in the test group the soft tissue apical to the root exposure was elevated full thickness by inserting a periosteum elevator in the probable sulcus. On the contrary, in the control group, a blade was used at the buccal aspect of the involved teeth, elevating the flap by a split thickness approach (Allen & Miller, 1989; da Silva et al., 2004). A statistically significant difference regarding the flap thickness in the two groups (0.93 ± 0.1 mm for test group and 0.46 ± 0.1 mm for control group) suggests that the “split–full–split” approach is able to preserve entirely the soft tissue thickness of the sulcular area.

Moreover, in this study, linear regression analysis showed a significant association between recession reduction and flap thickness resulted after the elevation. The critical role of soft tissue thickness for successful root coverage with the CAF technique was well reported in the previous literature. Baldi et al. (1999) demonstrated, in 19 patients that flap thickness and recession reduction are directly related, considering that a flap thickness >0.8 mm was associated with the possibility to obtain complete root coverage. The same association was evidenced also in a systematic review investigating flap thickness as a predictor of recession coverage (Hwang & Wang, 2006).

The successful results in terms of root coverage achieved in the test group of the present study were associated with a slight increase in KTH. These findings are in agreement with previously published data, describing single recessions treated with a “split–full–split” approach (Del Pizzo, Zucchelli, Modica, Villa, & Debernardi, 2005; Modica, Del Pizzo, Rocuzzo, & Romagnoli, 2000; de Sanctis & Zucchelli, 2007). The control group (PT) conversely displayed a decrease in the height of keratinized tissue. This clinical behaviour, observed when complete split thickness was performed, agrees as well with previous data (da Silva et al., 2004). Actually, in the latter investigation, 6 months after a complete “split” CAF, a slight loss of KT was observed.

At the best author’s knowledge, this is the first investigation that compares in the same population and in a randomized group two different techniques of CAF for the treatment of single recession defects. In a randomized controlled clinical trial, Mazzocco et al. (2011) investigated the efficacies of a partial- and full-thickness flap reflections, reporting no differences between study groups in terms of mean RC and KT gain after 6 months of healing. However, they treated multiple adjacent recessions by means of CAF with a subepithelial connective tissue graft (SCTG).

The Consensus Report of the Group 2 of the 10th Workshop on Periodontology (Tonetti & Jepsen, 2014) declared that patient-reported outcomes may represent the “true endpoints” of the mucogingival procedure, which capture patients’ perceptions of the therapy and complement conventional clinical outcomes. In this clinical trial, patient-centred outcomes were assessed by means of a VAS scale after 7 days (discomfort) and after 1 year (aesthetic and satisfaction). Moreover, the presence or absence of hypersensitivity was assessed.

Root sensitivity at baseline was detected in more than half of the patient population (test and control): The prevalence of this subjective symptom is consistent with that observed in other clinical trials (Pini Prato et al., 2005; Santamaria et al., 2017). After 1 year, root sensitivity was not detectable in the test group but it is still present, albeit in a small percentage of individuals, in the control group. This finding can be related with the greater surface of exposed root still present in the control group at the end of the experimental period and by the lower number of cases in which a CRC was obtained.

The discomfort perceived by individuals part of the control group after the first week of healing was twice as intense as that perceived by the test group. This interesting feature can be partially explained by the greater amount of blood clot produced by the more extensive partial thickness performed in the control group and its consequent greater difficulty to be reabsorbed in the very first phase of healing (first week).

After 12 months, VAS aesthetic and satisfaction were, respectively, 8.80 ± 1 and 8.47 ± 1.1 for the test group and 8.20 ± 1.6 and 8.20 ± 1.6 for the control group, confirming that clinical improvement in RC and the achievement of a CRC are important for the patient's perspective. Considering a range from 5 (neutral) to 10 (very good) of the VAS scale, a significant difference was found in favour of the test group for VAS satisfaction, probably due to the presence in the control group of a higher percentage of root sensitivity.

5 | CONCLUSIONS

Within the limits of this study, the following conclusion can be drawn:

- Thickness modulation of the split–full–split approach preserves soft tissue thickness of the flap at the maximum extent in the portion that will cover the denuded root.
- Flap thickness preservation and the presence of the periosteum in part of the flap may play a role in achieving better recession reduction outcomes (higher percentages of RC and CRC) at 1 year, with better patient's centred outcomes (higher hypersensitivity reduction at 1 year and lower VAS discomfort at 7 days)).
- The successful results in terms of root coverage achieved in the present study were not associated with a clinically significant increase in KTH at 1 year.

CONFLICT OF INTEREST

The authors report no conflict of interests related to the study.

ORCID

Marco Clementini  <http://orcid.org/0000-0002-1827-5200>

REFERENCES

- Allen, E. P., & Miller, P. D. (1989). Coronal positioning of existing gingiva: Short term results in the treatment of shallow marginal tissue recession. *Journal of Periodontology*, *60*, 316–319. <https://doi.org/10.1902/jop.1989.60.6.316>
- Baldi, C., Pini Prato, G. P., Pagliaro, U., Nieri, M., Saletta, D., Muzzi, L., & Cortellini, P. (1999). Coronally advanced flap procedure for root coverage. Is flap thickness a relevant predictor to achieve root coverage? A 19-case series. *Journal of Periodontology*, *70*, 1077–1084. <https://doi.org/10.1902/jop.1999.70.9.1077>
- Bernimoulin, J. P., Luscher, B., & Muhlemann, H. (1975). Coronally repositioned periodontal flap. *Journal of Clinical Periodontology*, *2*, 1–13. <https://doi.org/10.1111/j.1600-051X.1975.tb01721.x>
- Cairo, F., Nieri, M., & Pagliaro, U. (2014). Efficacy of periodontal plastic surgery procedures in the treatment of localized facial gingival recessions. A systematic review. *Journal of Clinical Periodontology*, *41*, S44–S62. <https://doi.org/10.1111/jcpe.12182>
- Chambrone, L., & Tatakis, D. N. (2015). Periodontal soft tissue root coverage procedures: A systematic review from the AAP Regeneration Workshop. *Journal of Periodontology*, *86*, 8–51. <https://doi.org/10.1902/jop.2015.130674>
- Chambrone, L., & Tatakis, D. N. (2016). Long-term outcomes of untreated buccal gingival recessions: A systematic review and meta-analysis. *Journal of Periodontology*, *87*, 796–808. <https://doi.org/10.1902/jop.2016.150625>
- Del Pizzo, M., Zucchelli, G., Modica, F., Villa, R., & Debernardi, C. (2005). Coronally advanced flap with or without enamel matrix derivative for root coverage: A 2-year study. *Journal of Clinical Periodontology*, *32*, 1181–1187. <https://doi.org/10.1111/j.1600-051X.2005.00831.x>
- Grieb, G., Steffens, G., Pallua, N., Bernhagen, J., & Bucala, R. (2011). Circulating fibrocytes- biology and mechanisms in wound healing and scar formation. *International Review of Cell and Molecular Biology*, *291*, 1–19.
- Haney, J. M., Nilveus, R. E., Mc Millan, P. J., & Wikesjö, U. M. (1993). Periodontal repair in dogs: Expanded polytetrafluoroethylene barrier membranes support wound stabilization and enhance bone regeneration. *Journal of Periodontology*, *64*, 883–890. <https://doi.org/10.1902/jop.1993.64.9.883>
- Hwang, D., & Wang, H. L. (2006). Flap thickness as a predictor of root coverage: A systematic review. *Journal of Periodontology*, *77*, 1625–1634. <https://doi.org/10.1902/jop.2006.060107>
- Kon, S., Novaes, A. B., Ruben, M. P., & Goldman, H. M. (1969). Visualization of the microvascularization of the healing periodontal wound. IV. Mucogingival surgery: Full thickness flap. *Journal of Periodontology* *40*, 441–456. <https://doi.org/10.1902/jop.1969.40.8.441>
- Laurens, N., Koolwijk, P., & de Maat, M. P. (2006). Fibrin structure and wound healing. *Journal of Thrombosis and Haemostasis*, *4*, 932–939. <https://doi.org/10.1111/j.1538-7836.2006.01861.x>
- Mazzocco, F., Comuzzi, L., Stefani, R., Milan, Y., Favero, G., & Stellini, E. (2011). Coronally advanced flap combined with a subepithelial connective tissue graft using full- or partial-thickness flap reflection. *Journal of Periodontology*, *82*, 1524–1529. <https://doi.org/10.1902/jop.2011.100586>
- Miller, P. D. (1985). A classification of marginal tissue recession. *The International Journal of Periodontics and Restorative Dentistry*, *5*, 8–13.
- Modica, F., Del Pizzo, M., Rocuzzo, M., & Romagnoli, R. (2000). Coronally advanced flap for the treatment of buccal gingival recessions with and without enamel matrix derivative. A split-mouth study. *Journal of Periodontology*, *71*, 1693–1698. <https://doi.org/10.1902/jop.2000.71.11.1693>
- Muhlemann, H. R., & Son, S. (1971). Gingival sulcus bleeding –a leading symptom in initial gingivitis. *Helvetica Odontologica Acta*, *15*, 107–113.
- Norberg, O. (1926). Ar en utlakning utan vovnad- sfortust otankbar vid kirurgisk behandling av. S. K. Alveolarpyorrhoe?. *Svensk Tandlaekare Tidsskrift* *19*, 171.

- O'Leary, T. J., Drake, R. B., & Naylor, J. E. (1972). The plaque control record. *Journal of Periodontology*, 43, 38. <https://doi.org/10.1902/jop.1972.43.1.38>
- Pini Prato, G. P., Baldi, C., Nieri, M., Franceschi, D., Cortellini, P., Clauser, C., ... Muzzi, L. (2005). Coronally advanced flap: The post-surgical position of the gingival margin is an important factor for achieving complete root coverage. *Journal of Periodontology*, 76, 713–722. <https://doi.org/10.1902/jop.2005.76.5.713>
- Pini Prato, G. P., Pagliaro, U., Baldi, C., Nieri, M., Saletta, D., Cairo, F., & Cortellini, P. (2000). Coronally advanced flap procedure for root coverage. Flap with tension versus flap without tension: A randomized controlled clinical study. *Journal of Periodontology*, 71, 188–201. <https://doi.org/10.1902/jop.2000.71.2.188>
- Pini Prato, G. P., Tinti, C., Vincenzi, G., Magnani, C., Cortellini, P., & Clauser, C. (1992). Guided tissue regeneration versus mucogingival surgery in the treatment of human buccal gingival recession. *Journal of Periodontology*, 63, 919–928. <https://doi.org/10.1902/jop.1992.63.11.919>
- Potente, M., Gerhardt, H., & Carmeliet, P. (2011). Basic and therapeutic aspects of angiogenesis. *Cell*, 146, 873–887. <https://doi.org/10.1016/j.cell.2011.08.039>
- de Sanctis, M., & Clementini, M. (2014). Flap approaches in plastic periodontal and implant surgery: Critical elements in design and execution. *Journal of Clinical Periodontology*, 41, 108–122. <https://doi.org/10.1111/jcpe.12189>
- de Sanctis, M., & Zucchelli, G. (2007). Coronally advanced flap: A modified surgical approach for isolated recession-type defects: Three-year results. *Journal of Clinical Periodontology*, 34, 262–268. <https://doi.org/10.1111/j.1600-051X.2006.01039.x>
- Santamaria, M. P., Neves, F. L. D. S., Silveira, C. A., Mathias, I. F., Fernandes-Dias, S. B., Jardini, M. A. N., & Tatakis, D. N. (2017). Connective tissue graft and tunnel or trapezoidal flap for the treatment of single maxillary gingival recessions: A randomized clinical trial. *Journal of Clinical Periodontology*, 44, 540–547. <https://doi.org/10.1111/jcpe.12714>
- da Silva, R. C., Joly, J. C., de Lima, A. F., & Tatakis, D. N. (2004). Root coverage using the coronally positioned flap with or without a subepithelial connective tissue graft. *Journal of Periodontology*, 75, 413–419. <https://doi.org/10.1902/jop.2004.75.3.413>
- Tonetti, M. S., Jepsen, S., & Working Group 2 of the European Workshop on Periodontology. (2014). Clinical efficacy of periodontal plastic surgery procedures: Consensus report of Group 2 of the 10th European Workshop on Periodontology. *Journal of Clinical Periodontology*, 41, S36–S43. <https://doi.org/10.1111/jcpe.12219>
- Wikesjö, U. M., Claffey, N., & Egelberg, J. (1991). Periodontal repair in dogs. Effect of heparin treatment of the root surface. *Journal of Clinical Periodontology*, 18, 60–64. <https://doi.org/10.1111/j.1600-051X.1991.tb01120.x>
- Wikesjö, U. M., Claffey, N., Nilvéus, R., & Egelberg, J. (1991). Periodontal repair in dogs: Effect of root surface treatment with stannous fluoride or citric acid on root resorption. *Journal of Periodontology*, 62, 180–184. <https://doi.org/10.1902/jop.1991.62.3.180>
- Wikesjö, U. M., & Nilvéus, R. (1990). Periodontal repair in dogs: Effect of wound stabilization on healing. *Journal of Periodontology*, 61, 719–724. <https://doi.org/10.1902/jop.1990.61.12.719>
- Zucchelli, G., Mele, M., Stefanini, M., Mazzotti, C., Marzadori, M., Montebugnoli, L., & de Sanctis, M. (2010). Patient morbidity and root coverage outcome after subepithelial connective tissue and de-epithelialized grafts: A comparative randomized-controlled clinical trial. *Journal of Clinical Periodontology*, 37, 728–738.

How to cite this article: Clementini M, Discepoli N, Danesi C, de Sanctis M. Biologically guided flap stability: The role of flap thickness including periosteum retention on the performance of the coronally advanced flap—A double-blind randomized clinical trial. *J Clin Periodontol*. 2018;45:1238–1246. <https://doi.org/10.1111/jcpe.12998>