

Time of morulation and trophectoderm quality are predictors of a live birth after euploid blastocyst transfer: a multicenter study

Laura Rienzi, M.Sc.,^a Danilo Cimadomo, Ph.D.,^a Arantxa Delgado, Ph.D.,^b Maria Giulia Minasi, M.Sc.,^c Gemma Fabozzi, M.Sc.,^a Raquel del Gallego, M.Sc.,^b Marta Stoppa, M.Sc.,^a Jose Bellver, M.D.,^b Adriano Gianciani, M.Sc.,^{a,d} Marga Esbert, Ph.D.,^e Antonio Capalbo, M.Sc.,^{d,f} Jose Remohi, M.D.,^b Ermanno Greco, M.D.,^c Filippo Maria Ubaldi, M.D., Ph.D.,^a and Marcos Meseguer, Ph.D.^b

^a Clinica Valle Giulia, G.EN.E.R.A. Centers for Reproductive Medicine, Rome, Italy; ^b IVI-RMA, Valencia, Spain; ^c Center for Reproductive Medicine, European Hospital, Rome, Italy; ^d DAHFMU, Unit of Histology and Medical Embryology, "Sapienza" University of Rome, Italy; ^e IVI-RMA, Barcelona, Spain; and ^f Igenomix, Marostica, Italy

Objective: To investigate whether the morphodynamic characterization of a euploid blastocyst's development allows a higher prediction of a live birth after single-embryo-transfer (SET).

Design: Observational cohort study conducted in two phases: training and validation.

Setting: Private in vitro fertilization centers.

Patient(s): Euploid blastocysts: 511 and 319 first vitrified-warmed SETs from 868 and 546 patients undergoing preimplantation genetic testing for aneuploidies (PGT-A) in the training and validation phase, respectively.

Intervention(s): Data collected from time of polar body extrusion to time of starting blastulation, and trophectoderm and inner-cell-mass static morphology in all embryos cultured in a specific time-lapse incubator with a continuous medium. Logistic regressions conducted to outline the variables showing a statistically significant association with live birth. In the validation phase, these variables were tested in an independent data set.

Main Outcome Measure(s): Live births per SET.

Result(s): The average live birth rate (LBR) in the training set was 40% (N = 207/511). Only time of morulation (tM) and trophectoderm quality were outlined as putative predictors of live birth at two IVF centers. In the validation set, the euploid blastocysts characterized by tM < 80 hours and high-quality trophectoderm resulted in a LBR of 55.2% (n = 37/67), while those with tM ≥ 80 hours and a low-quality trophectoderm resulted in a LBR of 25.5% (N = 13/51).

Conclusion(s): Time of morulation and trophectoderm quality are better predictors of a euploid blastocyst's reproductive competence. Our evidence was reproducible across different centers under specific culture conditions. These data support the crucial role of morulation for embryo development, a stage that involves massive morphologic, cellular, and molecular changes and deserves more investigation. (Fertil Steril® 2019;112:1080–93. ©2019 by American Society for Reproductive Medicine.)

El resumen está disponible en Español al final del artículo.

Key Words: Blastocyst, embryo selection, euploid, morula, time-lapse microscopy

Discuss: You can discuss this article with its authors and other readers at <https://www.fertsterdialog.com/users/16110-fertility-and-sterility/posts/51405-28408>

Received June 4, 2019; revised July 11, 2019; accepted July 22, 2019.

L.R. has received grants from Grant for Fertility Innovation – Merck, Forward 2016 – Finox, Investigator Sponsored Trial – Merck; personal fees for lectures from Merck, Merck-Serono, MSD, Excemed, and IBSA; partnership/stockholder from Genera Health Care srl and Flam srl, outside the submitted work. D.C. has received personal fees for lectures from Irvine Scientific, outside the submitted work. A.D. has nothing to disclose. M.G.M. has received personal fees for lectures from Cook Medical, outside the submitted work. G.F. has nothing to disclose. R.D. has nothing to disclose. M.S. has nothing to disclose. J.B. has nothing to disclose. A.G. has nothing to disclose. M.E. has nothing to disclose. A.C. has nothing to disclose. J.R. has nothing to disclose. E.G. has nothing to disclose. F.M.U. has received grants from Grant for Fertility Innovation – Merck, Forward 2016 – Finox, Investigator Sponsored Trial – Merck; personal fees for lectures from Merck, Merck-Serono, MSD, Excemed, and IBSA; partnership/stockholder from Genera Health Care SRL and Flam SRL, outside the submitted work. M.M. has received speaker fees from MSD, Merck and Ferring.

Reprint requests: Laura Rienzi, M.Sc., GENERA Centers for Reproductive Medicine, Rome, Naples, Marostica and Umbertide, Italy (E-mail: rienzi@generaroma.it).

Fertility and Sterility® Vol. 112, No. 6, December 2019 0015-0282/\$36.00

Copyright ©2019 American Society for Reproductive Medicine, Published by Elsevier Inc.

<https://doi.org/10.1016/j.fertnstert.2019.07.1322>

In vitro fertilization (IVF) today, achieving a high chance of live birth with concurrent low reproductive hazards per transfer (such as multiple pregnancy or miscarriage) is a task of utmost importance. To this end, some scientific societies have supported the implementation of a single-embryo transfer (SET) policy (1, 2) to limit the impact of multiple pregnancies (3). However, SET requires efficient embryo selection to be clinically sustainable (4).

Numerous studies have investigated the predictive value of parameters such as number of cells, their size and symmetry, and fragmentation for use in adopting a cleavage-stage embryo transfer policy (5). However, the results have been disappointing, probably because of the high impact of chromosomal aneuploidies in day 3 of preimplantation development (6), which indirectly hinders any other embryonic feature. Furthermore, natural selection during the first 3 days of culture is limited, so the cohort of cleavage-stage embryos produced in an IVF cycle is very similar to the initial cohort of fertilized oocytes (7, 8). The extension of embryo culture to the blastocyst stage generally reduces the cohort of embryos by 40% to 50% (7, 8). Thus, working in standard, controlled, constantly monitored culture conditions (9) might result in efficient selection in blastocyst culture. In fact, the data inspected by Glujovsky and Farquhar in 2016 in their Cochrane review (10, 11) showed improved implantation rates per embryo transfer when a blastocyst rather than a cleavage-stage embryo was transferred, with a similar cumulative live birth rate (LBR) (where the cumulative LBR is the overall live births achieved from consecutive fresh and frozen transfers of all the embryos produced in an IVF cycle). In other words, higher efficiency can be achieved without impacting the overall chance of success even when a single blastocyst is transferred, with a goal of limiting multiple gestations (3). Even so, a (single) blastocyst transfer policy does not reduce the rate of spontaneous miscarriages (11).

For an embryo cohort produced by a couple during an IVF cycle, an advanced strategy for selection involves retrieving about 5 to 10 trophoctoderm (TE) cells from each blastocyst and analyzing them via comprehensive chromosome testing (CCT) techniques to identify the euploid and aneuploid embryos. The aim of this approach—known as preimplantation genetic testing for aneuploidies (PGT-A) (12, 13)—is to prevent the transfer of aneuploid embryos. Dahdouh et al. (14) and Chen et al. (15) in two meta-analyses published in 2015 reported that when CCT is adopted to identify euploid embryos in good-prognosis patients, higher implantation rates per embryo transfer (~50–60%) and lower miscarriage rates per clinical pregnancy (~10%) can be achieved. This evidence encouraged more confident implementation of a blastocyst SET policy in women with a good prognosis and those of advanced maternal age (16, 17). In the latter population, higher efficiency with no impact on IVF efficacy (i.e., fewer embryo transfers, more implantations per embryo transfer, and lower miscarriage rates with similar cumulative LBR per patient with respect to non-PGT-A cycles) was reported from both the randomized controlled trials conducted by Ru-


bio and colleagues and the ESHRE Study into the Evaluation of Oocyte Euploidy by Microarray Analysis (ESTEEM). This was true of euploidy diagnosed by CCT on blastomeres and polar bodies (18, 19), which were less reliable specimens than TE for PGT-A purposes (20, 21).

Yet even if a normal chromosomal constitution is critical in predicting embryo implantation outcomes, it does not represent the only feature determining embryo competence—there is still room for improvement in embryo selection. Moreover, TE biopsy, albeit reported safe in the hands of experienced operators (21–25), still represents an invasive procedure that involves embryo manipulation and requires proper training and expertise. Therefore, the quest for noninvasive criteria is ongoing. Unfortunately, although several intriguing metabolomic/proteomic approaches have been adopted to analyze the byproducts of IVF (e.g., follicular fluid, cumulus cells, spent culture media), the clinical efficiency of these approaches has yet to be determined. Indeed, more extensive research and validation are needed before any consensus can be achieved or a clinical application developed (4, 26, 27).

The most commonly used method for embryo selection worldwide still is static morphology. At the blastocyst stage, numeric or alphabetic scales have been proposed (28), but all are based on the scoring system proposed by Gardner and Schoolcraft in 1999 (29), which consisted in the grading of three different variables: expansion, cohesiveness of the TE, and dimension and form of the inner cell mass (ICM). Several studies have attempted to identify the individual contribution of these parameters to implantation potential or live births, but different conclusions were reported. Some investigators emphasized the crucial role of the TE over the ICM (30–36); others instead emphasized the latter parameter as more predictive of a live birth (37–39).

The assessment of static morphology remains highly subjective (40), and most importantly static morphology is only mildly associated with chromosomal aneuploidies, which so far still represent the main cause of implantation failure and miscarriage in humans (41–45). Static morphologic evaluation has been improved with the introduction of time-lapse microscopy (TLM), which allows the assessment of embryo morphodynamic patterns throughout preimplantation development. The benefit of time-lapse incubators in providing an undisturbed culture environment is generally recognized, but TLM parameters have produced contrasting results when adopted to foresee embryo euploid constitution or reproductive competence. Their predictive value has not been consistent across studies and is still subject to debate (46–50). Importantly, morphokinetics appears to be affected by several parameters, such as infertility factor, maternal age, stimulation protocol, and culture strategies (51), which clearly limits the wide reproducibility of the algorithms used to date (52–55).

We propose that even if morphology (static and dynamic) only in part allows envisaging the chromosomal constitution

FIGURE 1

Flowchart of study phase 1 (training set to outline the variables associated with live birth after vitrified-warmed euploid blastocyst single embryo transfer) and phase 2 (validation set to define the consistency and reproducibility of those predictive variables). MII = metaphase II oocyte; PGT-A = preimplantation genetic testing of aneuploidies; SET = single-embryo transfer.

Rienzi. Euploid blastocysts' morphodynamic analysis. *Fertil Steril* 2019.

or the implantation competence of untested embryos, it still can be investigated for its potential to predict the latter in blastocysts diagnosed as euploid after PGT-A. To this end, we assessed the main variables associated with live birth after euploid blastocyst transfer in a data set from two IVF centers (training set). These variables were then tested in an independent data set that included a third IVF center (validation set). All the IVF centers involved in this study are reference laboratories for blastocyst culture, PGT, and vitrified-warmed SET. Our overarching aim was to address the consistency and reproducibility of any morphologic/morphodynamic parameter associated with live birth and identified in a clinical setting, subject to the lowest number of putative sources

of bias: undisturbed culture with a continuous media and euploid blastocyst SET.

MATERIALS AND METHODS

Study Design

The study design, mean maternal age, and main cycle data are shown in Figure 1. In the training phase, an observational cohort study of PGT-A cycles was performed between January 2016 and September 2017 at two European IVF centers: G.EN.E.R.A. Center for Reproductive Medicine (Rome, Italy; IVF center 1) and IVI (Valencia, Spain; IVF center 2). This phase was aimed at identifying the variables associated

with a live birth after the first vitrified-warmed euploid blastocyst SETs at both IVF centers.

In the validation phase, the variables outlined in the training set were tested for consistency and reproducibility on an independent data set. The latter set encompassed the first vitrified-warmed euploid blastocyst SETs performed after PGT-A between September 2017 and June 2018 at IVF centers 1 and 2 and at a third clinic (Center for Reproductive Medicine, European Hospital, Rome, Italy; IVF center 3).

Only the first SETs of euploid vitrified-warmed blastocysts were included in both study phases to avoid a putative cohort effect upon the LBR achieved in this population of patients. All PGT-A cycles entailing embryo culture conducted with a sequential media were not included for the sake of consistency within and between the centers, and because a lower blastulation and a slower development with respect to a continuous medium was previously reported at IVF center 1 (56).

IVF Procedures

Controlled ovarian stimulation was performed as previously described elsewhere (57). Thirty-five hours after ovulation trigger, the oocytes were collected by transvaginal ultrasound-guided aspiration. Fertilization was conducted only by intracytoplasmic sperm injection (ICSI) after denudation, as previously described elsewhere (58). Embryo culture was conducted individually in 25 μ L of continuous single-culture medium (CSCM; Irvine Scientific) covered by preequilibrated mineral oil in a microwell of the EmbryoSlide (Vitrolife). No media changeover in day 3 was performed.

Only an EmbryoScope time-lapse incubator (Vitrolife) was used at IVF centers 1, 2, and 3. Images of each embryo were acquired every 15 minutes on seven different focal planes. Embryos were cultured up to the blastocyst stage in 6% CO₂ and 5% O₂. In IVF centers 1 and 3, a simultaneous zona opening and TE biopsy approach (41) was used; IVF center 2 adopted a day-3 hatching-based TE biopsy approach (59). We conducted PGT-A by comprehensive chromosome testing techniques (60–64). The transferred blastocysts were selected independent from their morphodynamic parameters and based only on an euploid chromosomal constitution.

All euploid SETs were performed in a subsequent prepared cycle (57). The luteal phase was supported by vaginal micronized progesterone, 400 mg/day (Progeffik; Effik). We did not perform SET in cases of [1] an endometrium <7 mm and/or not trilaminar, [2] endometrial fluid in the uterine cavity, [3] hydrosalpinx, and/or [4] high levels of thyroid-stimulating hormone (≥ 2.5 IU/L). Vitrification (conducted within 30 minutes of the TE biopsy) and warming were performed according to previous publications (23, 45, 65).

Training phase. Between January 2016 and September 2017, 868 PGT-A cycles performed at IVF centers 1 and 2 were submitted to continuous embryo culture in an EmbryoScope incubator. The first vitrified-warmed euploid blastocyst SETs meeting the inclusion criteria were 147 and 364 from IVF centers 1 and 2, respectively (see Fig. 1). The time-lapse parameters adopted in this study were previously defined by Ciray et al. (66): timing of second polar body appearance

(tPB2); timing of completion of cleavage to 2, 3, 4, 5, and 8 cells (t2, t3, t4, t5, and t8, respectively); timing of morula formation (tM, the stage where visualization of individual blastomeres was no longer possible and full compaction was achieved); timing of starting blastocyst formation (tSB); length of the first (cc1 = t2 – tPB2), second (cc2 = t4 – t2), and third cell cycle (cc3 = t8 – t4); length of synchronization of cell divisions (s2 = t4 – t3); and synchronization of cleavage pattern (s3 = t8 – t5).

The time of blastocyst formation (tB) and blastocyst full expansion (tEB) were not included among the parameters under investigation because a different blastocyst biopsy approach was conducted at IVF center 2. Specifically, zona drilling in this center was performed on day 3, a procedure that induces a precocious hatching of the blastocyst (tHB) before the achievement of full expansion status (tEB) so that the TE biopsy and blastocyst grading are conducted significantly before with respect to IVF center 1, where both grading and zona drilling are instead routinely performed at the tEB.

Therefore, to reach a consensus between the different IVF centers involved in the study, the grading of the ICM and TE quality was here performed at the tB. The ICM and TE were graded according to the parameters defined in the Istanbul consensus (28, 67). Class 1 and class 2 were clustered together and defined as “high quality,” whereas class 3 was defined “low quality.” All the morphodynamic and static morphologic criteria were tested through logistic regression analyses adjusted for confounders.


Validation phase. Between September 2017 and June 2018, 546 PGT-A cycles performed at IVF centers 1, 2, and 3 were submitted to continuous embryo culture in an EmbryoScope incubator. The variables found to be statistically significantly associated with a live birth after euploid blastocyst SETs were tested for consistency and reproducibility on a further 319 vitrified-warmed euploid blastocyst SETs conducted at IVF centers 1 (N = 74) and 2 (N = 168) as well as in a third clinic (IVF center 3, N = 77) (see Fig. 1). The study was approved by the institutional review boards of the contributing clinics.

All the evaluations were performed by a single expert operator at each center. In a preliminary phase, the chosen operators were compared for their consistency in the definition of tM and in TE grading at tB on a set of 40 randomly chosen consecutive pictures (some examples can be found in Fig. 2). They showed an interclass correlation coefficient of 0.9 and 0.8 for the two variables, respectively. Both these values indicate an excellent reproducibility according to the thresholds defined by Koo and Li in 2016 (68).

Statistical Analysis

The investigated euploid blastocysts were divided into two groups based on whether they resulted in a live birth. Continuous variables are shown as mean $\pm$ standard deviation (SD) and range. Shapiro-Wilk tests were conducted to investigate whether the data followed a normal (Gaussian) distribution. Student's *t*-tests or Mann-Whitney *U* tests were conducted to assess statistically significant differences. Categorical variables are shown as rate with 95% confidence interval (CI).

FIGURE 2



Fisher's exact or chi-square tests were conducted to assess statistically significant differences.

We investigated the morphodynamic and static criteria from univariate and multivariate regression analyses by defining the binary response parameter as a live birth achieved ("1") or not ("0"). If required, the morphokinetic timings were converted from continuous variables into categorical by defining the cutoff value corresponding to the 50th percentile of prediction of a live birth. $P < .05$ was considered statistically significant. Statistical analyses were performed using the Statistical Package for the Social Sciences 19 (SPSS; IBM, Inc.). Power calculations were performed using G*Power v3.1 (Universität Düsseldorf).

RESULTS

Training Phase: Definition of the Variables Associated with a Live Birth after Vitrified-Warmed Euploid Blastocyst Transfer

In the training phase, 272 and 596 PGT-A cycles were conducted at IVF centers 1 and 2, respectively. The mean maternal age and main cycle data are shown in Figure 1. The cohort of metaphase 2 (MII) oocytes collected at IVF center 2 was larger than at IVF center 1 ($P < .01$), but then no difference was shown in terms of blastulation or euploidy rates. At IVF center 1, 49 (18%) and 76 (28%) PGT-A cycles resulted in no blastocyst and no euploid blastocyst produced, respectively. At IVF center 2, 98 (16%) and 134 (22.5%) PGT-A cycles resulted in no blastocyst and no euploid blastocyst produced, respectively. Therefore, 147 and 364 first vitrified-warmed euploid blastocyst SETs were included from IVF centers 1 and 2, respectively (see Fig. 1). The overall LBR per first vitrified-warmed euploid blastocyst SET was 40.5% ($N = 207/511$; 95% CI, 36.2–44.9). There was no difference between the two IVF centers (43.5%; 95% CI, 35.5–52.0;

$N = 64/147$ vs. 39.3%; 95% CI, 34.3–44.5; $N = 143/364$, respectively; $P = \text{not statistically significant [NS]}$).

Overall, the euploid embryos cultured at IVF center 2 rather than at IVF center 1 were consistently slower from tPB2 up to t5 (on average 1–2 hours). Statistically significant differences were indeed reported for tPB2, cc1, s2, cc2, and t5. However, after the eight-cell stage was reached, they could manage to decrease their delay; thus, starting from t8 onward no statistically significant difference was reported thereafter between the two centers (Supplemental Table 1, available online).

Of note, the tM was the first moment of preimplantation development where the timings at the two centers were almost perfectly overlapping (87.4 ± 9.2 , 66.9–116.0 hours vs. 87.4 ± 10.0 , 58.2–115.6 hours at centers 1 and 2, respectively). When the data in the two centers were clustered according to clinical outcome after first vitrified-warmed euploid blastocyst SETs (live birth vs. implantation failure/miscarriage), again different behaviors were reported during the stages before t8. Specifically, statistically significant differences were reported among reproductively competent and incompetent euploid blastocysts at center 2 for t2, t3, and t5, which were not confirmed at center 1 (Supplemental Table 2, available online). Conversely, the two centers were concordant for the differences reported for t8, cc3, s3, and tM, all statistically significantly longer in nonimplanted/miscarried euploid blastocysts (see Supplemental Table 2).

The tSB was also statistically significantly longer in nonimplanted/miscarried euploid blastocysts only at center 2 (see Supplemental Table 2). We did not include the blastulation timings any further (tB and tEB) due to the different blastocyst biopsy policies adopted at the two centers (i.e., a simultaneous zona opening and TE cell retrieval policy versus a day-3 zona opening-based approach at centers 1 and 2, respectively).

TABLE 1

Logistic regression analysis of the parameters associated with a live birth after euploid vitrified-warmed single-blastocyst transfer.

Parameter	Univariate			Multivariate		
	OR	95% CI	P value	OR	95% CI	P value
Maternal age	0.99	0.94–1.04	NS	—	—	—
IVF center						
1	—	—	—	—	—	—
2	0.84	0.57–1.23	NS	—	—	—
tPB2	0.86	0.74–0.99	.04	0.89	0.76–1.04	NS
tM	0.96	0.94–0.98	<.01	0.97	0.94–0.99	.04
tSB	0.97	0.95–0.99	<.01	1.03	0.99–1.07	NS
t2	0.95	0.89–1.00	NS	—	—	—
t3	0.96	0.93–1.00	NS	—	—	—
t4	0.95	0.92–0.99	.01	0.96	0.86–1.09	NS
t5	0.97	0.95–1.0	.05	1.05	0.95–1.17	NS
t8	0.95	0.93–0.97	<.01	0.97	0.93–1.02	NS
cc1	0.97	0.91–1.04	NS	—	—	—
cc2	0.96	0.91–1.01	NS	—	—	—
cc3	0.95	0.92–0.97	<.01	0.96	0.87–1.05	NS
s2	0.96	0.91–1.01	NS	—	—	—
s3	0.94	0.92–0.97	<.01	1.03	0.93–1.14	NS
ICM at tB						
High quality	—	—	—	—	—	—
Low quality	0.41	0.26–0.67	<.01	0.77	0.43–1.37	NS
TE at tB						
High quality	—	—	—	—	—	—
Low quality	0.30	0.19–0.45	<.01	0.38	0.23–0.63	<.01

Note: All embryos were cultured with the same continuous culture media and in the same time-lapse incubator in two different IVF centers. Static morphologic scoring was performed at tB (time of blastulation) in both centers. The Istanbul consensus criteria were adopted and class 1 and 2 were defined as high-quality, and class 3 was considered as low quality. Timing of second polar body appearance (tPB2); timing of cleavage to 2, 3, 4, 5, and 8 cells (t2, t3, t4, t5, and t8); timing of morula formation (tM); timing of starting blastocyst formation (tSB); timing of blastocyst formation (tB); length of the first, second and third cell cycle (cc1, cc2, and cc3); length of synchronization of cell divisions (s2) and synchronization of cleavage pattern (s3). CI = confidence interval; ICM = inner cell mass; NS = not statistically significant; OR = odds ratio; TE = trophectoderm.

Rienzi. Euploid blastocysts' morphodynamic analysis. *Fertil Steril* 2019.

At IVF center 1 a shorter period of time also was required to reach full expansion from euploid blastocysts, resulting in a live birth with respect to nonimplanted/miscarried ones (tEB – tSB: 18.0 ± 5.6 hours, 7.7–34.6 vs. 20.9 ± 7.5 hours, 4.5–43.3, respectively; $P=.01$; Supplemental Fig. 1, available online). In future studies, only centers adopting a simultaneous zona opening and trophectoderm biopsy approach would better assess these dynamics.

Table 1 shows the logistic regression analysis. Neither maternal age nor IVF center showed any association with live birth. Among the morphodynamic parameters under investigation, a statistically significant association was reported from the univariate analyses for tPB2, t4, t5, s3, cc3, t8, tM, and tSB. However, the tM was the only parameter found to be statistically significant from the multivariate analysis (odds ratio [OR] 0.97; 95% CI, 0.94–0.99; $P=.04$). Among the static morphologic features at tB, both the ICM and TE were statistically significantly associated with a live birth after euploid blastocyst vitrified-warmed SET from the univariate analyses. However, only the TE was still statistically significant from the multivariate analysis (OR 0.38; 95% CI, 0.23–0.63; $P<.01$).


A cutoff value for tM was then calculated based on the logistic regression analysis. In detail, a tM of 80 hours during the preimplantation development of an euploid blastocyst corresponded to the 50th percentile of prediction of a live birth after vitrified-warmed SET in the training set. Therefore, we clustered the results in four groups according to TE

morphologic quality evaluated at tB (high versus low) and tM (<80 hours versus ≥ 80 hours) (Fig. 3A). Interestingly, the euploid blastocysts characterized by a high-quality TE at tB and a tM <80 hours resulted in a LBR of 62.5% ($N = 45/72$; 95% CI, 50.3–73.4) after vitrified-warmed SET while the euploid blastocysts characterized by a low-quality TE at tB and a tM >80 hours resulted in a LBR of 21.0% ($N = 33/157$; 95% CI, 15.0–28.4) (see Fig. 3A).

Validation Phase: Definition of the Reproducibility of the Predictive Variables in an Independent Multicenter Data Set

The predictive power of the variables was tested on an independent data set from IVF centers 1 and 2 as well as from IVF center 3. This design allowed us to test the consistency and reproducibility on this validation set. In this study phase, 126, 280, and 140 PGT-A cycles were conducted at IVF center 1, 2, and 3, respectively. The mean maternal age and main cycle data are shown in Figure 1. Also in this case, the cohort of MII oocytes collected at IVF center 2 was larger than at IVF center 1 and 3 ($P=.02$), but then no difference was shown in terms of blastulation or euploidy rates.

After blastocyst culture and aneuploidy testing, 74, 168, and 77 first vitrified-warmed euploid blastocyst SETs could be included from IVF centers 1, 2, and 3, respectively. The rate of cycles where no blastocysts (16%, 14%, and 10%) and no euploid blastocysts (25%, 26%, and 35%) were

FIGURE 3

obtained were similar across the three clinics. The overall LBR per first vitrified-warmed euploid blastocyst SET was 42.9% (N = 137/319), consistent among the three IVF centers (45.9%, 40.5%, and 45.5% at IVF centers 1, 2, and 3, respectively).

Figure 3B shows the LBR in each specific group. For group A, we found the high-quality TE and tM <80 hours (N = 37/67, 55.2%; 95% CI, 42.6–67.2). For group B, we found the high-quality TE and tM ≥80 hours (N = 82/184, 44.6%; 95% CI, 37.3–52.1). For group C, we found the low-quality TE and tM <80 hours (N = 5/17, 29.4%; 95% CI, 11.4–55.9). For group D, we found the low-quality TE and tM ≥80 hours (N = 13/51, 25.5%; 95% CI, 14.8–39.9) (chi-squared=0.01; Fisher's exact test group A versus D, $P<.01$, and power = 0.9, group B versus D, $P=.02$ and power =

0.7). The results were consistent among the three IVF centers (e.g., group A: 61.5%, 55.0%, and 50.0% at IVF centers 1, 2, and 3, respectively; group D: 30.0%, 25.0%, and 23.1% at IVF centers 1, 2, and 3, respectively).

DISCUSSION

Over the past decade, morphokinetic analysis of embryo development has become an exciting area of human assisted reproduction. Expectations for it were high because of the amount of previously inaccessible information that could be finally collected. Even more importantly, the majority of data was either intrinsically quantitative (including time of well-defined events) or could be quantified (for example, some degenerative processes), signifying a considerable

opportunity to measure important features related to embryo competence.

Time-lapse technology has been therefore tested over these years to discern the relationship between morphokinetics and embryo developmental competence up to the blastocyst stage (55, 69–75) with different proposed algorithms (55, 76, 77), or between morphokinetics and implantation potential (53, 71, 76, 78–83) with the implementation of other algorithms (76, 79, 81). Also further associations were reported between some peculiar morphodynamic features (intercellular contact number or blastocyst collapse) and implantation potential (77, 84). Also specific dysmorphisms such as multinucleation, reverse cleavage, direct unequal cleavage, and irregular chaotic division have been revealed in human preimplantation embryos as related to embryo developmental or reproductive competence (85–89). Moreover, different trials were conducted to combine morphokinetics with proteomic and metabolomic investigations as well as with cumulus cells' gene expression patterns (90–92), aiming at improved predictivity. Future studies are needed to confirm the value such attempts in this field.

Thus, the controversial results for morphokinetics in predicting embryo implantation may be explained by various external and internal factors—differences in terminologies and parameters investigated, various culture conditions, or time-lapse systems used. However, far more uncertainty remains with aneuploidies. Studies dealing with the predictive value of morphokinetics for aneuploidy have contradicted each other, and they do not seem to have provided robust results. Apparently, morphodynamic behavior is in fact similar, especially between euploid embryos and embryos affected from a single chromosomal impairment. Multiple aneuploidies might instead have a greater impact on the timing of preimplantation development, but their incidence in viable blastocysts might also be lower, thereby limiting their predictive power (45, 47, 52, 93–97).

In summary, despite an enormous amount of studies and a meta-analysis reporting moderate evidence in favor of time-lapse microscopy (98), the superiority of morphokinetic assessment over traditional microscopic evaluation of morphology or PGT-A cannot be supported at present (99). Moreover, euploidy status does not exceed 50% to 60% of positive predictive value upon blastocyst reproductive competence after SET (14). Therefore, the purpose of our study was to combine blastocyst morphologic static and dynamic evaluation with euploidy.

A previous group had designed a similar study (100) and adopted morphokinetic parameters to visualize competent embryos among euploid ones. Their results were encouraging but were based on a limited sample size. Our study instead accounts for a large sample size from a multicenter perspective in both the training and validation set. This strategy allowed us to ensure the consistency and reproducibility of our evidence. Of note, the embryos cultured in sequential media were excluded from this data set because a significantly lower blastulation and slower development with respect to the continuous media was reported at IVF center 1 in a previous study (56). Therefore, both the training and validation phases

were conducted at two to three IVF centers but using the same time-lapse incubator, in the same conditions, and only with a continuous media.

The TE quality at tB was the only parameter deriving from a static morphologic evaluation and concordant among the two IVF centers as associated with euploid blastocysts' reproductive competence in the training set. This finding is consistent with the many previous studies that claimed its importance in both fresh (30, 32–34) and vitrified-warmed cycles (31, 35, 36). Also, blastocyst expansion and hatching status have been previously reported as predictive of implantation (39), but here they could not be assessed in the absence of bias at both centers. In fact, the practice of zona drilling on day 3 at IVF center 2 limited the reliability of such evaluation and required embryo morphologic grading to be conducted at tB (and not at tEB) also at IVF center 1 for the sake of homogeneity between the two data sets.

Nonetheless, TE quality and blastocyst expansion might be directly associated, as some investigators proposed previously. Specifically, a high-quality TE might involve a more efficient pumping of the ions, in turn prompting the accumulation of fluid in the blastocoel cavity (30). Therefore, the main contrasting results derive only from those studies claiming a higher importance of the ICM in the establishment of a pregnancy (37, 38). In our study, the ICM quality—once corrected for the other factors associated with euploid blastocyst implantation (including TE quality)—lost its significance. This higher importance of the TE in the process of implantation might be ascribed to its roles in this crucial phase: [1] it is the origin of the placenta and its annexes, [2] it promotes hatching and invasion of the endometrium (30, 101), and [3] it establishes the communication with the maternal immune system by secreting human chorionic gonadotropin (hCG) and inducing immunological tolerance (102, 103). In this regard, in the 1990s, a better quality TE was reported to secrete higher levels of hCG at an earlier time (104, 105). Likewise, more recently, gene expression profiled from TE cells outlines the overexpression of gene families involved in cell adhesion and cell communication to be associated with positive pregnancy outcomes (106, 107). All of this is evidence that supports our findings.

The tM was the only parameter derived from a morphodynamic evaluation of embryo preimplantation development that was concordant among the two IVF centers in the training set as associated with euploid blastocysts' reproductive competence. Consistent delayed development was reported for nonimplanted euploid blastocysts already at t8, which then accumulated (longer cc3 and s3) to culminate in an average 4 hours longer tM. Indeed, this last parameter in itself was sufficient to result in a statistically significant association with the live birth.

It is important to recognize that, despite morulation being a crucial step in preimplantation development (108), little attention has been paid to this stage in the literature. Compaction and subsequent morula development involve the flattening of the blastomeres and the consequent formation of junctions between them, followed by a massive redistribution of surface microvilli and other components of the plasma membrane (109). The embryo shows also a concurrent

increase in transcriptional and translational processes, possibly resulting from the full activation of the embryonic genome (which in humans normally is initiated at the 4- to 8-cell transition) as well as a marked change in the patterns of phospholipid synthesis (110). The cells lose their totipotency and get polarized radially. Across this axis, a differential division occurs, which generates two populations of cells with unequal distribution of organelles: outer polar cells with surface microvilli that will originate the TE and inner apolar cells with tight junctions containing basal nuclei, which will originate the ICM (111). Although the exact timings of the onset of such clusterization are unknown, it might be regulated via posttranslational modifications of specific proteins such as E-cadherins, the putative key effectors enabling a tighter adhesion between the cells (112–116).

When investigating clinical studies, Kramer et al. (117) reported in 2014 the importance of tM. Specifically, they found a significant association between the duration of compaction and an euploid chromosomal constitution in human embryos—yet this parameter was insufficient to provide results comparable to the actual CCT. In addition, both delayed (118) and precocious (109) compaction have been reported to be detrimental for embryo developmental potential and quality from other groups. A study by Mizobe et al. (119) outlined the timing of compaction to be associated with eventual pregnancy outcome, although this was not confirmed by other studies (73, 120).

Finally, also some studies conducted with conventional incubation and static morphologic grading systems highlighted the importance of morula assessment to foresee blastocyst quality and/or competence (121–125). All these findings together with our data should prompt future investigation of the morulation stage of development. More data focused on this issue will indeed be pivotal in understanding how euploid embryos acquire, maintain, or lose their reproductive competence during morulation.

To our knowledge, only one study has compared day-3 zona opening versus simultaneous zona opening and TE fragment retrieval blastocyst biopsy methods (126). In a patient-based randomized controlled trial by Zhao et al. (126), no difference was reported between the two groups, apart from a higher number of blastocysts cryopreserved when the latter biopsy approach was used. Nevertheless, it might be argued that artificial zona opening on day 3 could affect the developmental timing between the cleavage and the blastocyst stage. Here, by comparing three centers using two different biopsy approaches, we could instead indirectly report that the presence of a hole in the zona pellucida probably does not impair the timing up to morulation. Thus, if a difference in embryo morphokinetic development does result from the presence of an opening in the zona pellucida from day 3, it might affect the timings after blastulation rather than before (i.e., blastocyst expansion and hatching).

In the training phase of this study the interval of time between starting blastulation and blastocyst full expansion (tEB – tSB) at IVF center 1 was also different between euploid blastocysts resulting in a live birth rather than an implantation failure/miscarriage, but this timing could not be verified at IVF center 2. Future single-center or multicenter studies

conducted solely in IVF laboratories adopting a simultaneous zona opening and trophectoderm biopsy approach are highly recommended to better address these dynamics and their predictive power upon euploid blastocyst implantation.

The main strengths of our study are [1] the definition of two variables statistically significantly associated with euploid blastocyst reproductive competence, whose consistency and reproducibility was independently validated in three IVF centers (euploid blastocysts with high-quality TE and tM <80 hours resulted in 62% and 55% LBR in the training and validation sets versus 21% and 25% for euploid blastocysts with low-quality TE and tM ≥80 hours); and [2] the applicability of these predictive variables regardless of the TE biopsy protocol adopted in each laboratory. Nevertheless, our study was conducted under specific culture conditions at the three centers (continuous media, same time lapse incubator and low oxygen tension atmosphere). Further investigations are needed to understand the strength of this association in different conditions.

Ideal future studies would also entail [1] a powered randomized controlled trial comparing a study arm where the first euploid vitrified-warmed blastocyst to transfer is chosen based on both TE quality and tM versus a control arm where no selection scheme is adopted, and [2] an investigation into the predictive power of these two variables during standard ICSI cycles in the absence of aneuploidy testing. Clearly the same conditions as in this report should be adopted for both these studies.

From a clinical perspective, this study highlights the importance of a rigorous validation and of analogous culture conditions when static and morphodynamic parameters are tested as selection criteria. In this regard, artificial intelligence and machine learning represent a concrete future perspective to further increase the standardization and objectivity of each evaluation performed by different operators in different clinics (127, 128).

From a biological perspective, the morula stage emerged as a putative crucial step for the acquisition/maintenance of reproductive competence across human embryo preimplantation development. Future studies will better unravel the cellular and molecular events characterizing the process of morulation.

REFERENCES

1. Practice Committee, Society for Assisted Reproductive Technology and the American Society for Reproductive Medicine. Guidelines on the number of embryos transferred. *Fertil Steril* 2004;82:773–4.
2. Harbottle S, Hughes C, Cutting R, Roberts S, Brison D. Association of Clinical Embryologists & the (ACE) British Fertility Society (BFS). Elective single embryo transfer: an update to UK Best Practice Guidelines. *Hum Fertil (Camb)* 2015;18:165–83.
3. Pandian Z, Marjoribanks J, Ozturk O, Serour G, Bhattacharya S. Number of embryos for transfer following in vitro fertilisation or intra-cytoplasmic sperm injection. *Cochrane Database Syst Rev* 2013;7:CD003416.
4. Bolton VN, Leary C, Harbottle S, Cutting R, Harper JC. How should we choose the 'best' embryo? A commentary on behalf of the British Fertility Society and the Association of Clinical Embryologists. *Hum Fertil (Camb)* 2015;18:156–64.
5. Machtinger R, Racowsky C. Morphological systems of human embryo assessment and clinical evidence. *Reprod Biomed Online* 2013;26:210–21.

6. McCoy RC, Newnham LJ, Ottolini CS, Hoffmann ER, Chatzimeletiou K, Cornejo OE, et al. Tripolar chromosome segregation drives the association between maternal genotype at variants spanning PLK4 and aneuploidy in human preimplantation embryos. *Hum Mol Genet* 2018;27:2573–85.
7. Stoop D, Van Landuyt L, Van den Abbeel E, Camus M, Verheyen G, Devroey P. Should a single blastocyst transfer policy be a clinical decision or should it depend on the embryological evaluation on day 3? *Reprod Biol Endocrinol* 2011;9:60.
8. Sfountouris IA, Kolibianakis EM, Lainas GT, Venetis CA, Petsas GK, Tarlatzis BC, et al. Blastocyst utilization rates after continuous culture in two commercial single-step media: a prospective randomized study with sibling oocytes. *J Assist Reprod Genet* 2017;34:1377–83.
9. Wale PL, Gardner DK. The effects of chemical and physical factors on mammalian embryo culture and their importance for the practice of assisted human reproduction. *Hum Reprod Update* 2016;22:2–22.
10. Glujovsky D, Farquhar C. Cleavage-stage or blastocyst transfer: what are the benefits and harms? *Fertil Steril* 2016;106:244–50.
11. Glujovsky D, Farquhar C, Quinteiro Retamar AM, Alvarez Sedo CR, Blake D. Cleavage stage versus blastocyst stage embryo transfer in assisted reproductive technology. *Cochrane Database Syst Rev* 2016;6:CD002118.
12. Zegers-Hochschild F, Adamson GD, Dyer S, Racowsky C, de Mouzon J, Sokol R, et al. The international glossary on infertility and fertility care, 2017. *Fertil Steril* 2017;108:393–406.
13. Zegers-Hochschild F, Adamson GD, Dyer S, Racowsky C, de Mouzon J, Sokol R, et al. The international glossary on infertility and fertility care, 2017. *Hum Reprod* 2017;32:1786–801.
14. Dahdouh EM, Balayla J, Garcia-Velasco JA. Comprehensive chromosome screening improves embryo selection: a meta-analysis. *Fertil Steril* 2015;104:1503–12.
15. Chen M, Wei S, Hu J, Quan S. Can comprehensive chromosome screening technology improve IVF/ICSI outcomes? A meta-analysis. *PLoS One* 2015;10:e0140779.
16. Forman EJ, Hong KH, Ferry KM, Tao X, Taylor D, Levy B, et al. In vitro fertilization with single euploid blastocyst transfer: a randomized controlled trial. *Fertil Steril* 2013;100:100–7.e1.
17. Ubaldi FM, Capalbo A, Colamaria S, Ferrero S, Maggiulli R, Vajta G, et al. Reduction of multiple pregnancies in the advanced maternal age population after implementation of an elective single embryo transfer policy coupled with enhanced embryo selection: pre- and post-intervention study. *Hum Reprod* 2015;30:2097–106.
18. Rubio C, Bellver J, Rodrigo L, Castillon G, Guillen A, Vidal C, et al. In vitro fertilization with preimplantation genetic diagnosis for aneuploidies in advanced maternal age: a randomized, controlled study. *Fertil Steril* 2017;107:1122–9.
19. Verpoest W, Staessen C, Bossuyt PM, Goossens V, Altarescu G, Bonduelle M, et al. Preimplantation genetic testing for aneuploidy by microarray analysis of polar bodies in advanced maternal age: a randomized clinical trial. *Hum Reprod* 2018.
20. Capalbo A, Romanelli V, Cimadomo D, Girardi L, Stoppa M, Dovere L, et al. Implementing PGD/PGD-A in IVF clinics: considerations for the best laboratory approach and management. *J Assist Reprod Genet* 2016;33:1279–86.
21. Cimadomo D, Capalbo A, Ubaldi FM, Scarica C, Palagiano A, Canipari R, et al. The impact of biopsy on human embryo developmental potential during preimplantation genetic diagnosis. *Biomed Res Int* 2016;2016:7193075.
22. Scott RT Jr, Upham KM, Forman EJ, Zhao T, Treff NR. Cleavage-stage biopsy significantly impairs human embryonic implantation potential while blastocyst biopsy does not: a randomized and paired clinical trial. *Fertil Steril* 2013;100:624–30.
23. Cimadomo D, Capalbo A, Levi-Setti PE, Soscia D, Orlando G, Albani E, et al. Associations of blastocyst features, trophectoderm biopsy and other laboratory practice with post-warming behavior and implantation. *Hum Reprod* 2018;33:1992–2001.
24. Cimadomo D, Rienzi L, Romanelli V, Alviggi E, Levi-Setti PE, Albani E, et al. Inconclusive chromosomal assessment after blastocyst biopsy: prevalence, causative factors and outcomes after re-biopsy and re-vitrification: a multicenter experience. *Hum Reprod* 2018;33:1839–46.
25. Capalbo A, Ubaldi FM, Cimadomo D, Maggiulli R, Patassini C, Dusi L, et al. Consistent and reproducible outcomes of blastocyst biopsy and aneuploidy screening across different biopsy practitioners: a multicentre study involving 2586 embryo biopsies. *Hum Reprod* 2016;31:199–208.
26. Gardner DK, Meseguer M, Rubio C, Treff NR. Diagnosis of human preimplantation embryo viability. *Hum Reprod Update* 2015;21:727–47.
27. Siristatidis CS, Sertedaki E, Vaidakis D, Varounis C, Trivella M. Metabolomics for improving pregnancy outcomes in women undergoing assisted reproductive technologies. *Cochrane Database Syst Rev* 2018;3:CD011872.
28. Alpha Scientists in Reproductive Medicine and ESHRE Special Interest Group of Embryology. The Istanbul consensus workshop on embryo assessment: proceedings of an expert meeting. *Hum Reprod* 2011;26:1270–83.
29. Gardner DK, Schoolcraft B. In vitro culture of human blastocyst. In: Jansen R, Mortimer D, editors. *Towards reproductive certainty: infertility and genetics beyond*. Carnforth, United Kingdom: Parthenon Press; 1999:377–88.
30. Ahlstrom A, Westin C, Reimer E, Wikland M, Hardarson T. Trophectoderm morphology: an important parameter for predicting live birth after single blastocyst transfer. *Hum Reprod* 2011;26:3289–96.
31. Ahlstrom A, Westin C, Wikland M, Hardarson T. Prediction of live birth in frozen-thawed single blastocyst transfer cycles by pre-freeze and post-thaw morphology. *Hum Reprod* 2013;28:1199–209.
32. Hill MJ, Richter KS, Heitmann RJ, Graham JR, Tucker MJ, DeCherney AH, et al. Trophectoderm grade predicts outcomes of single-blastocyst transfers. *Fertil Steril* 2013;99:1283–9.e1.
33. Thompson SM, Onwubalili N, Brown K, Jindal SK, McGovern PG. Blastocyst expansion score and trophectoderm morphology strongly predict successful clinical pregnancy and live birth following elective single embryo blastocyst transfer (eSET): a national study. *J Assist Reprod Genet* 2013;30:1577–81.
34. Ebner T, Tritscher K, Mayer RB, Oppelt P, Duba HC, Maurer M, et al. Quantitative and qualitative trophectoderm grading allows for prediction of live birth and gender. *J Assist Reprod Genet* 2016;33:49–57.
35. Chen X, Zhang J, Wu X, Cao S, Zhou L, Wang Y, et al. Trophectoderm morphology predicts outcomes of pregnancy in vitrified-warmed single-blastocyst transfer cycle in a Chinese population. *J Assist Reprod Genet* 2014;31:1475–81.
36. Honnma H, Baba T, Sasaki M, Hashiba Y, Ohno H, Fukunaga T, et al. Trophectoderm morphology significantly affects the rates of ongoing pregnancy and miscarriage in frozen-thawed single-blastocyst transfer cycle in vitro fertilization. *Fertil Steril* 2012;98:361–7.
37. Licciardi F, McCaffrey C, Oh C, Schmidt-Sarosi C, McCulloh DH. Birth weight is associated with inner cell mass grade of blastocysts. *Fertil Steril* 2015;103:382–7.e2.
38. Subira J, Craig J, Turner K, Bevan A, Ohuma E, McVeigh E, et al. Grade of the inner cell mass, but not trophectoderm, predicts live birth in fresh blastocyst single transfers. *Hum Fertil (Camb)* 2016;19:254–61.
39. Van den Abbeel E, Balaban B, Ziebe S, Lundin K, Cuesta MJ, Klein BM, et al. Association between blastocyst morphology and outcome of single-blastocyst transfer. *Reprod Biomed Online* 2013;27:353–61.
40. Storr A, Venetis CA, Cooke S, Kilani S, Ledger W. Inter-observer and intra-observer agreement between embryologists during selection of a single day 5 embryo for transfer: a multicenter study. *Hum Reprod* 2017;32:307–14.
41. Capalbo A, Rienzi L, Cimadomo D, Maggiulli R, Elliott T, Wright G, et al. Correlation between standard blastocyst morphology, euploidy and implantation: an observational study in two centers involving 956 screened blastocysts. *Hum Reprod* 2014;29:1173–81.
42. Alfawati S, Fragouli E, Colls P, Stevens J, Gutierrez-Mateo C, Schoolcraft WB, et al. The relationship between blastocyst morphology, chromosomal abnormality, and embryo gender. *Fertil Steril* 2011;95:520–4.

43. Heffner LJ. Advanced maternal age—how old is too old? *N Engl J Med* 2004;351:1927–9.
44. Hassold T, Hunt P. To err (meiotically) is human: the genesis of human aneuploidy. *Nat Rev Genet* 2001;2:280–91.
45. Minasi MG, Colasante A, Riccio T, Ruberti A, Casciani V, Scarselli F, et al. Correlation between aneuploidy, standard morphology evaluation and morphokinetic development in 1730 biopsied blastocysts: a consecutive case series study. *Hum Reprod* 2016;31:2245–54.
46. Kaser DJ, Racowsky C. Clinical outcomes following selection of human preimplantation embryos with time-lapse monitoring: a systematic review. *Hum Reprod Update* 2014;20:617–31.
47. Pennetta F, Lagalla C, Borini A. Embryo morphokinetic characteristics and euploidy. *Curr Opin Obstet Gynecol* 2018;30:185–96.
48. Basile N, Caiazza M, Meseguer M. What does morphokinetics add to embryo selection and in-vitro fertilization outcomes? *Curr Opin Obstet Gynecol* 2015;27:193–200.
49. Castello D, Motato Y, Basile N, Remohi J, Espejo-Catena M, Meseguer M. How much have we learned from time-lapse in clinical IVF? *Mol Hum Reprod* 2016;22:719–27.
50. Aparicio-Ruiz B, Romany L, Meseguer M. Selection of preimplantation embryos using time-lapse microscopy in in vitro fertilization: state of the technology and future directions. *Birth Defects Res* 2018;110:648–53.
51. Barrie A, Homburg R, McDowell G, Brown J, Kingsland C, Troup S. Examining the efficacy of six published time-lapse imaging embryo selection algorithms to predict implantation to demonstrate the need for the development of specific, in-house morphokinetic selection algorithms. *Fertil Steril* 2017;107:613–21.
52. Campbell A, Fishel S, Bowman N, Duffy S, Sedler M, Hickman CF. Modeling a risk classification of aneuploidy in human embryos using non-invasive morphokinetics. *Reprod Biomed Online* 2013;26:477–85.
53. Basile N, Vime P, Florensa M, Aparicio Ruiz B, Garcia Velasco JA, Remohi J, et al. The use of morphokinetics as a predictor of implantation: a multicentric study to define and validate an algorithm for embryo selection. *Hum Reprod* 2015;30:276–83.
54. Azzarello A, Hoest T, Mikkelsen AL. The impact of pronuclei morphology and dynamics on live birth outcome after time-lapse culture. *Hum Reprod* 2012;27:2649–57.
55. Cruz M, Garrido N, Herrero J, Perez-Cano I, Munoz M, Meseguer M. Timing of cell division in human cleavage-stage embryos is linked with blastocyst formation and quality. *Reprod Biomed Online* 2012;25:371–81.
56. Cimadomo D, Scarica C, Maggiulli R, Orlando G, Soscia D, Albrici L, et al. Continuous embryo culture elicits higher blastulation but similar cumulative delivery rates than sequential: a large prospective study. *J Assist Reprod Genet* 2018;35:1329–38.
57. Vaiarelli A, Cimadomo D, Patrizio P, Venturella R, Orlando G, Soscia D, et al. Biochemical pregnancy loss after frozen embryo transfer seems independent of embryo developmental stage and chromosomal status. *Reprod Biomed Online* 2018;37:349–57.
58. Rienzi L, Ubaldi F, Anniballo R, Cerulo G, Greco E. Preincubation of human oocytes may improve fertilization and embryo quality after intracytoplasmic sperm injection. *Hum Reprod* 1998;13:1014–9.
59. McArthur SJ, Leigh D, Marshall JT, de Boer KA, Jansen RP. Pregnancies and live births after trophoctoderm biopsy and preimplantation genetic testing of human blastocysts. *Fertil Steril* 2005;84:1628–36.
60. Treff NR, Tao X, Ferry KM, Su J, Taylor D, Scott RT Jr. Development and validation of an accurate quantitative real-time polymerase chain reaction-based assay for human blastocyst comprehensive chromosomal aneuploidy screening. *Fertil Steril* 2012;97:819–24.
61. Capalbo A, Treff NR, Cimadomo D, Tao X, Upham K, Ubaldi FM, et al. Comparison of array comparative genomic hybridization and quantitative real-time PCR-based aneuploidy screening of blastocyst biopsies. *Eur J Hum Genet* 2015;23:901–6.
62. Vera-Rodriguez M, Diez-Juan A, Jimenez-Almazan J, Martinez S, Navarro R, Peinado V, et al. Origin and composition of cell-free DNA in spent medium from human embryo culture during preimplantation development. *Hum Reprod* 2018;33:745–56.
63. Mir P, Rodrigo L, Mercader A, Buendia P, Mateu E, Milan-Sanchez M, et al. False positive rate of an arrayCGH platform for single-cell preimplantation genetic screening and subsequent clinical application on day-3. *J Assist Reprod Genet* 2013;30:143–9.
64. Fiorentino F, Biricik A, Bono S, Spizzichino L, Cotroneo E, Cottone G, et al. Development and validation of a next-generation sequencing-based protocol for 24-chromosome aneuploidy screening of embryos. *Fertil Steril* 2014;101:1375–82.
65. Cobo A, Bellver J, Domingo J, Perez S, Crespo J, Pellicer A, et al. New options in assisted reproduction technology: the Cryotop method of oocyte vitrification. *Reprod Biomed Online* 2008;17:68–72.
66. Ciray HN, Campbell A, Agerholm IE, Aguilar J, Chamayou S, Esbert M, et al. Proposed guidelines on the nomenclature and annotation of dynamic human embryo monitoring by a time-lapse user group. *Hum Reprod* 2014;29:2650–60.
67. ALPHA Scientists in Reproductive Medicine; ESHRE Special Interest Group Embryology. Istanbul consensus workshop on embryo assessment: proceedings of an expert meeting. *Reprod Biomed Online* 2011;22:632–46.
68. Koo TK, Li MY. A guideline of selecting and reporting intraclass correlation coefficients for reliability research. *J Chiropr Med* 2016;15:155–63.
69. Wong CC, Loewke KE, Bossert NL, Behr B, De Jonge CJ, Baer TM, et al. Non-invasive imaging of human embryos before embryonic genome activation predicts development to the blastocyst stage. *Nat Biotechnol* 2010;28:1115–21.
70. Conaghan J, Chen AA, Willman SP, Ivani K, Chenette PE, Boostanfar R, et al. Improving embryo selection using a computer-automated time-lapse image analysis test plus day 3 morphology: results from a prospective multicenter trial. *Fertil Steril* 2013;100:412–9.e5.
71. VerMilyea MD, Tan L, Anthony JT, Conaghan J, Ivani K, Gvakharina M, et al. Computer-automated time-lapse analysis results correlate with embryo implantation and clinical pregnancy: a blinded, multi-centre study. *Reprod Biomed Online* 2014;29:729–36.
72. Aparicio-Ruiz B, Basile N, Perez Albala S, Bronet F, Remohi J, Meseguer M. Automatic time-lapse instrument is superior to single-point morphology observation for selecting viable embryos: retrospective study in oocyte donation. *Fertil Steril* 2016;106:1379–85.e10.
73. Kirkegaard K, Kesmodel US, Hindkjaer JJ, Ingerslev HJ. Time-lapse parameters as predictors of blastocyst development and pregnancy outcome in embryos from good prognosis patients: a prospective cohort study. *Hum Reprod* 2013;28:2643–51.
74. Cetinkaya M, Pirkevi C, Yelke H, Colakoglu YK, Atayurt Z, Kahraman S. Relative kinetic expressions defining cleavage synchronicity are better predictors of blastocyst formation and quality than absolute time points. *J Assist Reprod Genet* 2015;32:27–35.
75. Milewski R, Kuc P, Kuczynska A, Stankiewicz B, Lukaszuk K, Kuczynski W. A predictive model for blastocyst formation based on morphokinetic parameters in time-lapse monitoring of embryo development. *J Assist Reprod Genet* 2015;32:571–9.
76. Motato Y, de los Santos MJ, Escriba MJ, Ruiz BA, Remohi J, Meseguer M. Morphokinetic analysis and embryonic prediction for blastocyst formation through an integrated time-lapse system. *Fertil Steril* 2016;105:376–84.e9.
77. Yang ST, Shi JX, Gong F, Zhang SP, Lu CF, Tan K, et al. Cleavage pattern predicts developmental potential of day 3 human embryos produced by IVF. *Reprod Biomed Online* 2015;30:625–34.
78. Kirkegaard K, Ahlstrom A, Ingerslev HJ, Hardarson T. Choosing the best embryo by time lapse versus standard morphology. *Fertil Steril* 2015;103:323–32.
79. Meseguer M, Herrero J, Tejera A, Hilligsoe KM, Ramsing NB, Remohi J. The use of morphokinetics as a predictor of embryo implantation. *Hum Reprod* 2011;26:2658–71.
80. Lemmen JG, Agerholm I, Ziebe S. Kinetic markers of human embryo quality using time-lapse recordings of IVF/ICSI-fertilized oocytes. *Reprod Biomed Online* 2008;17:385–91.
81. Liu Y, Chapple V, Feenan K, Roberts P, Matson P. Time-lapse deselection model for human day 3 in vitro fertilization embryos: the combination of

- qualitative and quantitative measures of embryo growth. *Fertil Steril* 2016; 105:656–62.e1.
82. Milewski R, Kuczynska A, Stankiewicz B, Kuczynski W. How much information about embryo implantation potential is included in morphokinetic data? A prediction model based on artificial neural networks and principal component analysis. *Adv Med Sci* 2017;62:202–6.
83. Milewski R, Milewska AJ, Kuczynska A, Stankiewicz B, Kuczynski W. Do morphokinetic data sets inform pregnancy potential? *J Assist Reprod Genet* 2016;33:357–65.
84. Liu Y, Chapple V, Feenan K, Roberts P, Matson P. Clinical significance of intercellular contact at the four-cell stage of human embryos, and the use of abnormal cleavage patterns to identify embryos with low implantation potential: a time-lapse study. *Fertil Steril* 2015;103:1485–91.e1.
85. Desai N, Ploskonka S, Goodman LR, Austin C, Goldberg J, Falcone T. Analysis of embryo morphokinetics, multinucleation and cleavage anomalies using continuous time-lapse monitoring in blastocyst transfer cycles. *Reprod Biol Endocrinol* 2014;12:54.
86. Athayde Wirka K, Chen AA, Conaghan J, Ivani K, Gvakharina M, Behr B, et al. Atypical embryo phenotypes identified by time-lapse microscopy: high prevalence and association with embryo development. *Fertil Steril* 2014;101:1637–48.e1–e5.
87. Meriano J, Clark C, Cadesky K, Laskin CA. Binucleated and micronucleated blastomeres in embryos derived from human assisted reproduction cycles. *Reprod Biomed Online* 2004;9:511–20.
88. Rubio I, Kuhlmann R, Agerholm I, Kirk J, Herrero J, Escriba MJ, et al. Limited implantation success of direct-cleaved human zygotes: a time-lapse study. *Fertil Steril* 2012;98:1458–63.
89. Zhan Q, Ye Z, Clarke R, Rosenwaks Z, Zaninovic N. Direct unequal cleavages: embryo developmental competence, genetic constitution and clinical outcome. *PLoS One* 2016;11:e0166398.
90. Hammond ER, Stewart B, Peek JC, Shelling AN, Cree LM. Assessing embryo quality by combining non-invasive markers: early time-lapse parameters reflect gene expression in associated cumulus cells. *Hum Reprod* 2015; 30:1850–60.
91. Dominguez F, Meseguer M, Aparicio-Ruiz B, Piqueras P, Quinonero A, Simon C. New strategy for diagnosing embryo implantation potential by combining proteomics and time-lapse technologies. *Fertil Steril* 2015; 104:908–14.
92. Scarica C, Cimadomo D, Dovere L, Giancani A, Stoppa M, Capalbo A, et al. An integrated investigation of oocyte developmental competence: expression of key genes in human cumulus cells, morphokinetics of early divisions, blastulation, and euploidy. *J Assist Reprod Genet* 2019;36:875–87.
93. Basile N, Nogales Mdel C, Bronet F, Florensa M, Riqueiros M, Rodrigo L, et al. Increasing the probability of selecting chromosomally normal embryos by time-lapse morphokinetics analysis. *Fertil Steril* 2014;101: 699–704.
94. Rienzi L, Capalbo A, Stoppa M, Romano S, Maggillulli R, Albricci L, et al. No evidence of association between blastocyst aneuploidy and morphokinetic assessment in a selected population of poor-prognosis patients: a longitudinal cohort study. *Reprod Biomed Online* 2015;30:57–66.
95. Chavez SL, Loewke KE, Han J, Moussavi F, Colls P, Munne S, et al. Dynamic blastomere behaviour reflects human embryo ploidy by the four-cell stage. *Nat Commun* 2012;3:1251.
96. Vera-Rodriguez M, Chavez SL, Rubio C, Reijo Pera RA, Simon C. Prediction model for aneuploidy in early human embryo development revealed by single-cell analysis. *Nat Commun* 2015;6:7601.
97. Patel DV, Shah PB, Kotdawala AP, Herrero J, Rubio I, Banker MR. Morphokinetic behavior of euploid and aneuploid embryos analyzed by time-lapse in embryoscope. *J Hum Reprod Sci* 2016;9:112–8.
98. Pribenszky C, Nilselid AM, Montag M. Time-lapse culture with morphokinetic embryo selection improves pregnancy and live birth chances and reduces early pregnancy loss: a meta-analysis. *Reprod Biomed Online* 2017; 35:511–20.
99. Armstrong S, Bhide P, Jordan V, Pacey A, Farquhar C. Time-lapse systems for embryo incubation and assessment in assisted reproduction. *Cochrane Database Syst Rev* 2018:CD011320.
100. Yang Z, Zhang J, Salem SA, Liu X, Kuang Y, Salem RD, et al. Selection of competent blastocysts for transfer by combining time-lapse monitoring and array CGH testing for patients undergoing preimplantation genetic screening: a prospective study with sibling oocytes. *BMC Med Genomics* 2014;7:38.
101. Norwitz ER, Schust DJ, Fisher SJ. Implantation and the survival of early pregnancy. *N Engl J Med* 2001;345:1400–8.
102. Licht P, Russu V, Wildt L. On the role of human chorionic gonadotropin (hCG) in the embryo-endometrial microenvironment: implications for differentiation and implantation. *Semin Reprod Med* 2001;19:37–47.
103. Tsampalas M, Griselet V, Berndt S, Foidart JM, Geenen V, Perrier d'Hauterive S. Human chorionic gonadotropin: a hormone with immunological and angiogenic properties. *J Reprod Immunol* 2010;85:93–8.
104. Dokras A, Sargent IL, Barlow DH. Human blastocyst grading: an indicator of developmental potential? *Hum Reprod* 1993;8:2119–27.
105. Lopata A. Implantation of the human embryo. *Hum Reprod* 1996;11(Suppl 1):175–84, discussion 93–95.
106. Jones GM, Cram DS, Song B, Kokkali G, Pantos K, Trounson AO. Novel strategy with potential to identify developmentally competent IVF blastocysts. *Hum Reprod* 2008;23:1748–59.
107. Parks JC, McCallie BR, Janesch AM, Schoolcraft WB, Katz-Jaffe MG. Blastocyst gene expression correlates with implantation potential. *Fertil Steril* 2011;95:1367–72.
108. Coticchio G, Lagalla C, Sturmeier R, Pennetta F, Borini A. The enigmatic morula: mechanisms of development, cell fate determination, self-correction and implications for ART. *Hum Reprod Update* 2019;25:422–38.
109. Iwata K, Yumoto K, Sugishima M, Mizoguchi C, Kai Y, Iba Y, et al. Analysis of compaction initiation in human embryos by using time-lapse cinematography. *J Assist Reprod Genet* 2014;31:421–6.
110. Braude P, Bolton V, Moore S. Human gene expression first occurs between the four- and eight-cell stages of preimplantation development. *Nature* 1988;332:459–61.
111. Elder K, Dale B. In-vitro fertilization. 3rd ed. Cambridge: Cambridge University Press; 2011.
112. Nikas G, Ao A, Winston RM, Handyside AH. Compaction and surface polarity in the human embryo in vitro. *Biol Reprod* 1996;55:32–7.
113. Fleming TP, Sheth B, Fesenko I. Cell adhesion in the preimplantation mammalian embryo and its role in trophectoderm differentiation and blastocyst morphogenesis. *Front Biosci* 2001;6:D1000–7.
114. Larue L, Ohsugi M, Hirchenhain J, Kemler R. E-cadherin null mutant embryos fail to form a trophectoderm epithelium. *Proc Natl Acad Sci USA* 1994;91:8263–7.
115. Bell CE, Calder MD, Watson AJ. Genomic RNA profiling and the programme controlling preimplantation mammalian development. *Mol Hum Reprod* 2008;14:691–701.
116. Alikani M. Epithelial cadherin distribution in abnormal human preimplantation embryos. *Hum Reprod* 2005;20:3369–75.
117. Kramer YG, Kofinas JD, Melzer K, Noyes N, McCaffrey C, Buldo-Licciardi J, et al. Assessing morphokinetic parameters via time lapse microscopy (TLM) to predict euploidy: are aneuploidy risk classification models universal? *J Assist Reprod Genet* 2014;31:1231–42.
118. Ivec M, Kovacic B, Vlaisavljevic V. Prediction of human blastocyst development from morulas with delayed and/or incomplete compaction. *Fertil Steril* 2011;96:1473–8.e2.
119. Mizobe Y, Ezono Y, Tokunaga M, Oya N, Iwakiri R, Yoshida N, et al. Selection of human blastocysts with a high implantation potential based on timely compaction. *J Assist Reprod Genet* 2017;34:991–7.
120. Chamayou S, Patrizio P, Storaci G, Tomaselli V, Alecci C, Ragolia C, et al. The use of morphokinetic parameters to select all embryos with full capacity to implant. *J Assist Reprod Genet* 2013;30:703–10.
121. Ebner T, Moser M, Shebl O, Sommergruber M, Gaiswinkler U, Tews G. Morphological analysis at compacting stage is a valuable prognostic tool for ICSI patients. *Reprod Biomed Online* 2009;18:61–6.
122. Fabozzi G, Alteri A, Rega E, Starita MF, Piscitelli C, Giannini P, et al. Morphological assessment on day 4 and its prognostic power in selecting viable embryos for transfer. *Zygote* 2016;24:477–84.
123. Tao J, Tamis R, Fink K, Williams B, Nelson-White T, Craig R. The neglected morula/compact stage embryo transfer. *Hum Reprod* 2002;17:1513–8.

124. Feil D, Henshaw RC, Lane M. Day 4 embryo selection is equal to day 5 using a new embryo scoring system validated in single embryo transfers. *Hum Reprod* 2008;23:1505–10.
125. Skiadas CC, Jackson KV, Racowsky C. Early compaction on day 3 may be associated with increased implantation potential. *Fertil Steril* 2006;86:1386–91.
126. Zhao H, Tao W, Li M, Liu H, Wu K, Ma S. Comparison of two protocols of blastocyst biopsy submitted to preimplantation genetic testing for aneuploidies: a randomized controlled trial. *Arch Gynecol Obstet* 2019;299:1487–93.
127. Khosravi P, Kazemi E, Zhan Q, Malmsten JE, Toschi M, Zisimopoulos P, et al. Deep learning enables robust assessment and selection of human blastocysts after in vitro fertilization. *NPJ Digital Med* 2019;2:21.
128. Tran D, Cooke S, Illingworth PJ, Gardner DK. Deep learning as a predictive tool for fetal heart pregnancy following time-lapse incubation and blastocyst transfer. *Hum Reprod* 2019;34:1011–8.

El tiempo de morulación y la calidad del trofoectodermo son predictores de nacido vivo tras la transferencia de un blastocisto euploide: un estudio multicéntrico

Objetivo: Investigar si la caracterización morfodinámica del desarrollo de un blastocisto euploide permite una mayor predicción de un nacido vivo después de la transferencia de un solo embrión (SET).

Diseño: Estudio de cohorte observacional realizado en dos fases: entrenamiento y validación.

Entorno: Centros privados de fertilización in vitro.

Paciente(s): Blastocistos euploides: 511 y 319 primeros SET vitrificados y descongelados de 868 y 546 pacientes sometidos a cribado genético preimplantacional de aneuploidías (PGT-A) en la fase de entrenamiento y validación, respectivamente.

Intervención(es): Recopilación de datos desde el momento de la extrusión del corpúsculo polar hasta el momento de comenzar la blastulación, y la morfología estática del trofoectodermo y la masa celular interna en embriones en una incubadora *time-lapse* con un medio continuo. Realización de regresiones logísticas para mostrar la asociación estadísticamente significativa con el nacido vivo. En la fase de validación, estas variables se probaron en un conjunto de datos independiente.

Principales medidas de resultados: Nacimientos vivos por SET

Resultados: La tasa promedio de nacido vivo (LBR) en el grupo de entrenamiento fue del 40% (N = 207/511). Solo el tiempo de morulación (tM) y la calidad del trofoectodermo se mostraron como supuestos predictores en dos de los centros de FIV. En el grupo de validación, los blastocistos euploides caracterizados por tM <80 horas y trofoectodermo de alta calidad resultaron en un LBR de 55.2% (n = 37/67), mientras que aquellos con tM ≥ 80 horas y un trofoectodermo de baja calidad resultaron en un LBR de 25.5% (N = 13/51).

Conclusión(es): El tiempo de morulación y la calidad del trofoectodermo son mejores predictores de la competencia reproductiva del blastocisto euploide. Nuestra evidencia ha sido reproducible en diferentes centros bajo condiciones de cultivo específicas. Estos datos respaldan el papel crucial de la morulación para el desarrollo embrionario, una etapa que implementa cambios morfológicos, celulares y moleculares masivos y merece más investigación.

SUPPLEMENTAL FIGURE 1

